

## Adjusting to a new home: Mycobacterium tuberculosis gene expression in response to an intracellular lifestyle.

Article (Accepted Version)

Stokes, Richard W and Waddell, Simon J (2009) Adjusting to a new home: Mycobacterium tuberculosis gene expression in response to an intracellular lifestyle. *Future Microbiology*, 4 (10). pp. 1317-1335. ISSN 1746-0921

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/7636/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

"Adjusting to a new home - *Mycobacterium tuberculosis* gene expression in response to an intracellular lifestyle."

Richard W. Stokes PhD<sup>1</sup> & Simon J. Waddell PhD<sup>2</sup>

<sup>1</sup> Department of Paediatrics, University of British Columbia, & Division of Infectious and Immunological Diseases, British Columbia's Children's Hospital, Child and Family Research Institute, 950 West 28<sup>th</sup> Avenue, Vancouver, British Columbia, Canada, V5Z 4H4.

Email: [rstokes@interchange.ubc.ca](mailto:rstokes@interchange.ubc.ca) Phone: 604 875 2471 Fax 604 875 2221

<sup>2</sup> Medical Microbiology, Centre for Infection, Division of Cellular & Molecular Medicine, St. George's University of London, Cranmer Terrace, Tooting, London, SW17 0RE, UK

Email: [swaddell@sgul.ac.uk](mailto:swaddell@sgul.ac.uk) Phone: 020 8725 2679 Fax: 020 8672 0234

## SUMMARY

*Mycobacterium tuberculosis* is still the most significant single species of bacteria causing disease in mankind. The ability of *M. tuberculosis* to survive and replicate within host macrophages is a pivotal step in its pathogenesis. Understanding the microenvironments that *M. tuberculosis* encounters within the macrophage and the adaptations that the bacterium undergoes to facilitate its survival will lead to insights into possible therapeutic targets for improved treatment of tuberculosis. This is urgently needed with the emergence of multi- and extensively-drug resistant strains of *M. tuberculosis*. Significant advances have been made in understanding the macrophage response on encountering *M. tuberculosis*. Complementary information is also accumulating regarding the counter responses of *M. tuberculosis* during the various stages of its interactions with the host. As such, a picture is emerging delineating the gene expression of intracellular *M. tuberculosis* at different stages of the interaction with macrophages.

## KEYWORDS

*Mycobacterium tuberculosis*; macrophage; microarray; host-pathogen interaction; phagosome; gene expression; innate immunity; adaptive immunity

## INTRODUCTION

*Mycobacterium tuberculosis* (*M.tb*) is the etiological agent of tuberculosis (TB) and is the leading cause of infectious death in adults due to a single bacterial species<sup>1</sup>. It is estimated that every year there are approximately 8 million new cases of TB, and that TB is the cause of around 2-3 million deaths worldwide<sup>301</sup>. Synergy with the human immunodeficiency virus (HIV), along with the emergence of multidrug resistant TB (MDR) and the newly identified extensively drug resistant (XDR) strains<sup>2</sup> threatens to make the disease incurable once again.

*M.tb* is a facultative intracellular pathogen possessing mechanisms for evading and subverting host immune responses. Only 10% of people infected with *M.tb* develop clinical disease over their lifetime. Thus, there is obviously a struggle between pathogen and host, with the outcome of any infection being greatly affected by variation in host susceptibility<sup>3 4 5 6</sup> and mycobacterial virulence<sup>7 8</sup>. That pathogenic mycobacteria are able to survive within macrophages (MM) and granulomas (both components of the host defence system) demonstrates that they have mechanisms enabling them to inhibit, endure or evade cell-mediated immunity. It is logical to propose that specific mycobacterial genes are critical for the survival and virulence of *M.tb* within the host. This hypothesis has led to numerous studies, using a wide range of methodologies, to recognise these functionally significant genes.

1  
2 The production of transposon mutant libraries in mycobacteria<sup>9 10</sup> has enabled the study of a  
3 number of potential virulence factors. Signature tagged mutagenesis has been used to identify  
4 mutants that do not survive *in vivo*<sup>11 12</sup>, whereas saturation mutagenesis of *M.tb* has identified  
5 mutants that can not survive *in vivo*<sup>13</sup> or in MM<sup>14</sup>. While informative, these studies have  
6 identified only the genes that are essential for the survival of *M.tb* in MM and animal models.  
7 Obviously this is important information that can be used in the search for new therapeutics but it  
8 does not highlight conditionally essential genes required in additional phases of infection, or  
9 allow for the identification of non-essential genes that mediate the differences seen when  
10 comparing, for instance, virulent and avirulent strains.

11  
12 Gene expression profiling on a global level has been investigated widely using such methods  
13 as microarray analysis, subtractive hybridisation and differential display. Differential display of  
14 RNA from two populations of bacteria (e.g. broth grown versus inside a MM) is a sensitive  
15 method that requires very small amounts of RNA. However, relatively large numbers of  
16 candidate bands must be checked to find a few differential bands and rare mRNA transcripts are  
17 very hard to detect without nested priming<sup>15</sup>. Subtractive hybridization has been used to  
18 eliminate bacterial genes commonly expressed in both MM and broth culture<sup>16 17 18</sup>. The  
19 remaining genes are then assumed to be specifically expressed in one or the other environment.  
20 Another approach to identifying genes expressed in MM is to construct a library of mycobacterial  
21 genomic DNA in a promoter capture vector, with the DNA inserted upstream of a promoterless  
22 Green Fluorescent Protein (GFP) gene. Thus, only plasmids containing an active promoter will  
23 express GFP. This library is transformed into the mycobacteria of interest which are then used to  
24 infect MMs. Any promoter constructs that are more actively expressed in the MMs can be  
25 identified using fluorescent readout methodologies<sup>19 20 21</sup>. Subtractive hybridization and  
26 promoter capture have been used successfully to identify genes expressed by mycobacteria within  
27 MMs. However, both methods have drawbacks in that they are technically demanding and are  
28 prone to identifying genes that can be found in both pathogenic and non-pathogenic  
29 mycobacteria.

30  
31 Post-genomic studies offer the potential for identifying putative virulence genes by utilising  
32 microarray technology to define the changing pattern of *M.tb* gene expression during infection<sup>22</sup>.  
33 DNA sequences representing the open reading frames (ORF) for all genes from (a) reference  
34 genome(s) are printed or synthesised on a glass slide (or chip) which is then probed with cDNA  
35 derived from *M.tb* RNA. The fluorophore-labelled cDNA is co-hybridised in a competitive  
36 reaction against cDNA (labelled with an alternative fluorophore) derived from a different  
37 condition in a direct comparison, or against a DNA reference that allows multi-way comparisons  
38 between RNA samples. The normalized ratio of the fluorophores for each gene reflects the  
39 change in RNA abundance between the two conditions. Developments such as the use of  
40 genome-directed primers<sup>23</sup>, bacterial RNA amplification<sup>24</sup>, and high-density microarray  
41 platforms have expanded the range of infection models tractable to investigation and the global  
42 nature of the data acquired. The application of transcriptome sequencing using next-generation  
43 methodologies will clearly improve the genomic resolution of these studies further.

44  
45 Another approach for studying transcriptomes is the use of Bacterial Artificial Chromosome  
46 (BAC) arrays. They have previously been used to compare mycobacterial genomes<sup>25 26 27 28</sup> and,  
47 more recently to compare the transcriptomes of intracellular *M.tb* strains<sup>29 30</sup>. Genomic DNA is  
48 cloned into BACs and the clones are positioned on a restriction enzyme map of the genome to  
49 produce a set of BACs that cover the genome. Southern blots of BAC digests can be hybridized  
50 with cDNA from two transcriptomes and non hybridizing fragments in one or the other sample  
51 can be identified. A drawback to this method is that each band produced by the restriction digest  
52 of each BAC does not necessarily represent a single gene. The absence of a gene within a band  
53 may, therefore be masked by a positive hybridisation with other genes within that same band.

Conversely, the presence of multiple genes in a band can be advantageous in identifying differences in the expression of operons<sup>30</sup>.

#### OVERVIEW OF THE INFECTIOUS PROCESS

*M.tb* bacilli first encounter host MMs following their inhalation into the alveolar spaces of the lung. It is here where they will encounter the alveolar MM. The first stage of the interaction is the attachment of the bacteria to the MM, followed by ingestion via phagocytosis (see Fig 1). During this process, the bacterium may be exposed to potentially toxic agents such as reactive oxygen intermediates (ROI). They will also trigger the production of cytokines and chemokines by the MM which play pivotal roles in the induction of innate immunity, adaptive immunity and apoptosis. Following ingestion the bacterium resides within an intracellular compartment called a phagosome where they are potentially exposed to additional killing mechanisms such as lysosomal hydrolases, antimicrobial peptides and glutathione. In addition, they must acquire nutrients while residing in this intracellular compartment of the MM usually associated with digestion and recycling of senescent cellular debris. Another important adaptation the TB bacillus must make is the move to a lower oxygen tension within the MM. Obviously some bacteria survive this initial move to the intracellular environment where they can then replicate. One outcome of this replication is the release of bacteria from the initial contact cell to other local host cells such as dendritic cells and newly arriving elicited MMs. Another outcome of bacterial replication is that the host develops adaptive immunity leading to the interferon- $\gamma$  (IFN $\gamma$ )-mediated activation of infected MMs. Activated MMs are better able to control bacterial replication through enhanced killing mechanisms such as the production of ROI and reactive nitrogen intermediates (RNI), phagosomal maturation to lysosomes, and apoptosis. The exposure to adaptive immune responses triggers a shift of the bacterial physiology towards a so called dormant or chronic phase. In this state, *M.tb* utilises lipids as a carbon source and the glyoxalate shunt for energy production. The host response to the persisting bacteria is to form a highly organised granuloma that walls off the bacteria in an hypoxic environment where they are commonly found within foamy macrophages. This granuloma may calcify over time or may cavitate and break through to the airways, releasing bacteria to begin the cycle again.

There are a number of critical steps during the cycle of *M.tb* pathogenesis that result in changes to the environment in which the bacterium finds itself. Many of these key phases of infection involve changes to the M $\Phi$ -bacterial interplay. Although *M.tb* has been shown to enter neutrophils<sup>31 32</sup> and epithelial cells<sup>33 34</sup>, the MM is considered to be the host cell in which the bacteria most commonly reside. While the role of neutrophils and epithelial cells in the pathogenesis of *M.tb* is still uncertain, the central role of the MM is generally accepted as a major component in the host/pathogen interaction in TB. Thus, this review will concentrate on the expression of genes by *M.tb* during its interaction with MMs. Additionally, although numerous informative studies have identified expression differences between broth grown *M.tb* and bacteria obtained from the human lung and animal infection models, or by using model organisms like *M. bovis* BCG<sup>35 36 37 38 39 40</sup>, we have restricted our review to studies utilising *M.tb* in the context of life inside the MM.

#### BINDING AND UPTAKE (MOVEMENT FROM EXTRACELLULAR TO INTRACELLULAR)

Because mycobacteria are predominantly found within MMs and are known to survive and replicate within that niche, it is generally believed that they have evolved strategies that facilitate their association with phagocytes. It is often assumed that mycobacteria readily enter any M $\Phi$  they encounter and it is only then that survival mechanisms of bacterial origin come into play. However, there is evidence that M $\Phi$  populations vary in their ability to efficiently associate with *M.tb*<sup>41 42</sup>. Thus, evidence is accumulating to suggest that pathogenic mycobacteria are not automatically bound and ingested by the first M $\Phi$  they encounter on inhalation. On the contrary, *M.tb* bacilli appear to have developed strategies to control their ingestion by MMs, which includes a capsule that acts to regulate the receptor-ligand interactions between *M.tb* and MMs<sup>43</sup>.

Host-derived serum opsonins can greatly enhance the uptake of mycobacteria by most MΦ populations<sup>44 45 42</sup>, though not by resident alveolar MMs<sup>42</sup>. The serum-independent interaction of *M.tb* with MMs is considered to be important in the early stages of pulmonary infection because the lung is an environment in which serum opsonins are relatively scarce or absent<sup>46 47</sup> and alveolar MΦs have low levels of receptors for serum opsonins<sup>48 42</sup>. Serum-mediated uptake of *M.tb* may be more important at later stages of the infection, when bacteria encounter elicited MΦs<sup>41 42</sup>. Thus, in the absence of significant levels of serum (a situation encountered by *M.tb* on entering the lung) *M.tb* appears to have developed strategies to limit its association with MΦs and to direct its entry into certain sub-populations of MΦs. Entry of *M.tb* into the MΦ appears to rely on bacterial adherence to MΦ receptors/surface molecules followed by phagocytosis, with no substantial evidence for active mechanisms of bacterial invasion. Numerous receptors have been implicated in mediating the serum-independent entry of *M.tb* into MΦs including the mannose receptor<sup>49</sup>, the lipopolysaccharide receptor (CD14),<sup>50</sup> surfactant protein receptors A<sup>51</sup>, the complement receptor CR3<sup>52</sup>, and CD43<sup>53 54</sup>.

It might be expected that binding and ingestion of *M.tb* to MΦ would trigger the induction of a number of genes to prepare the bacterium for the major changes in oxygen tension, nutrient availability and exposure to toxic molecules. Surprisingly, the attachment of *M.tb* to MΦ did not alter bacterial gene expression, as measured by microarray with a >1.5x fold change in expression compared to *M.tb* bacilli resuspended in MΦ culture media alone<sup>55</sup>. However, uptake of the bacteria over 2 hours resulted in transcriptional responses commencing within minutes and resulting in distinct changes in gene subsets over time<sup>55</sup>. These included members of the WhiB family of transcriptional regulators, members of the *devR* (also known as *dosR*) regulon involved in the response to hypoxia and ROI and >50% of the *phoPR* regulon, including genes involved in cell wall remodelling.

Several groups have defined the transcriptome of intracellular *M.tb* from 4h to 48h after MΦ infection, describing the changes in gene expression between broth grown and intracellular *M.tb*. Many additional studies have investigated specific parameters considered to be representative of the intracellular environment, such as hypoxia or low pH, and have been used to deconvolute the complex patterns of gene expression observed after macrophage uptake. Figure 2 illustrates the overlap of gene subsets between the intracellular transcriptional signatures derived from MΦ infection models and microarray experiments designed to mimic specific environmental stresses that *M.tb* may encounter during infection. Interestingly, there is limited overlap between all intracellular transcriptomes (A-D). However, common functionalities of regulated gene sets are repeatedly identified (E-S). These include genes indicative of fatty acid metabolism, cell wall metabolism and restructuring, iron metabolism, DNA damage repair, nutrient deprivation, transcriptional regulation and a response to hypoxia including the *devR* regulon<sup>56 57 58 59 55 60</sup>. These differences are taken to imply that the bacterium has moved to an environment low in carbohydrate yet relatively lipid rich, where it encounters a low oxygen tension, and is exposed to potentially toxic mechanisms requiring restructuring of the cell wall and the utilization of defence mechanisms. The specific events following uptake of *M.tb* by MΦ have been modelled and investigated in some detail.

#### RESIDENCE IN A PHAGOSOME AND THE ASSOCIATED LOW NUTRIENTS, INCREASED ACIDITY AND HYPOXIA.

The bacterial phagosome is a defining feature of *M.tb* pathogenesis as it differs from most particulate endosomes. There are numerous studies in which this compartment has been modelled in a cell free system. Following the phagocytosis of most particles, the phagosome will mature along the phagosomal-lysosomal pathway, sequentially acquiring markers from early endosomes (such as transferrin receptors, mannose receptors and Rab5), late endosomes (such as LAMPs, cathepsin D and Rab7) and lysosomes (such as LAMPs, and cathepsin D) via a “kiss

1 and run” phenomenon<sup>61 62</sup>. The phagosome also progressively acidifies down to pH5.5, or  
2 lower, following fusion with late endosomes and lysosomes<sup>63 64</sup>. However, pathogenic  
3 mycobacteria such as *M.tb* subvert this phagosome maturation, presumably to aid in their  
4 survival within the MΦ. Seminal work by Hart first demonstrated that mycobacterial  
5 phagosomes did not fuse with lysosomes<sup>65</sup>. This observation was extended to show that *M.tb*  
6 arrests maturation of the phagosome to reside in an early endosomal-like compartment that can  
7 interact with other endosomes, thereby obtaining nutrients for its survival. The *M.tb* phagosome  
8 contains the early endosome markers glycosphingolipids<sup>66</sup>, transferrin<sup>67</sup> and Rab5 but excludes  
9 the late endosome marker Rab7<sup>68</sup>. However, the late endosomal-lysosomal markers LAMP-1  
10 and cathepsin D do appear to be found in the *M.tb* phagosome<sup>69 70</sup> (Fig 1).  
11

12 The mycobacterial phagosome remains at pH 6.2-6.3 displaying a reduced acidification when  
13 compared to those containing control particles<sup>71 72</sup>. This has been attributed to a paucity of  
14 proton ATPase pumps on the *M.tb* phagosome<sup>69 73</sup>. Evidence has been presented to explain how  
15 mycobacteria bring about the reduced acidification and arrest of maturation of their phagosome.  
16 Galectin 3<sup>74</sup>, coronin1 (TACO)<sup>75</sup> and degraded cellubrevin<sup>76</sup> have all been found to accumulate  
17 on the *M.tb* phagosome but, as yet, no mechanism for their activity has been fully defined. The  
18 mycobacterial cell wall associated glycolipids, lipoarabinomannan<sup>77 78</sup> and cord factor<sup>79</sup> have  
19 been shown to inhibit phagosome maturation, though again, no mechanistic explanation has been  
20 determined. Inhibition of intracellular calcium by mycobacteria has also been shown to inhibit  
21 phagosome maturation<sup>80</sup> through the inhibition of sphingosine kinase<sup>81</sup>. In addition, transposon  
22 mutants of *M.tb* that fail to prevent the maturation of phagosome development have been  
23 identified and the genes involved were found to include putative transporters, lipid synthesis  
24 enzymes<sup>82</sup> and several mutations in the ESX-1 secretion pathway<sup>83</sup>. The importance of the  
25 proteins secreted by the ESX-1 pathway in down-regulating MΦ responses and in the virulence  
26 of *M.tb* has been recently reviewed<sup>84</sup>. That progression of endosomal maturation is deleterious to  
27 the survival of *M.tb* has been suggested by numerous studies<sup>82 85 86 87</sup>, though it is difficult to  
28 differentiate whether the death of intracellular *M.tb* is the result of, or precedes the development  
29 of, phagosome maturation.  
30

31 Exposure of *M.tb* to various models of the phagosome has resulted in a better understanding  
32 of the *M.tb* transcriptional response to this niche. Although the *M.tb* phagosome is held at a  
33 modified early endosome stage, thereby allowing access to some extracellular nutrients, there is  
34 still a level of nutrient deprivation the bacterium has to contend with, including low levels of  
35 iron. Growth of *M.tb* in a chemostat under nutrient depletion resulted in the bacteria entering a  
36 non-replicative state and an up-regulation of genes involved in  $\beta$  oxidation of fatty acids<sup>88</sup>.  
37 Starvation of *M.tb* led to a downregulation of 15% of the energy metabolism genes, 47% of the  
38 aerobic respiration genes, 59% of the genes involved in ribosomal protein synthesis and 16% of  
39 the lipid biosynthesis genes<sup>89</sup>. Interestingly, protein analysis identified significant upregulation  
40 of the *hspx* gene product,  $\alpha$ -crystallin that is associated with a dormancy profile<sup>89</sup>. Iron  
41 starvation leads to upregulation of genes involved in iron acquisition (*mbtA*, *mbtB*, *mbtI*)<sup>90</sup> and  
42 the regulation of a variety of other proteins, including putative transporters, iron storage proteins,  
43 members of the PE/PPE family, transcriptional regulators, and enzymes involved in lipid  
44 metabolism<sup>91</sup>. Many of these transcriptional signatures are mirrored in the intracellular gene  
45 expression patterns of *M.tb* after MM infection<sup>56 55 58 60 92</sup>.  
46

The regulation of genes following pH reduction is dependent on the relative acidity that *M.tb* is exposed to. The mild acidification (pH 6.2-6.3) seen in the “normal” *M.tb* phagosome results in the regulation of 162 genes, whereas exposure to a pH 5.5, that may be seen in IFN $\gamma$ -activated MMs, results in the differential regulation of 679 genes<sup>55</sup>. In another study in which *M.tb* was exposed to a pH of 5.5, 81 genes were differentially expressed with many of the highly induced genes showing homology with nonribosomal peptide synthetases/polyketide synthases<sup>93</sup>. Using a promoter trap system, it was shown that acid induced the expression of *lipF* (a putative lipase) and Rv0834c, though neither was induced in resting M $\Phi$  phagosomes<sup>94</sup>. Several members of the *phoP* regulon were induced by acid pH as was *whiB3* suggesting a role for these genes in regulation of the acid response<sup>55</sup>. A number of genes were only regulated by pH 5.5 and not pH6.5; these included repression of genes mediating protein synthesis, replication and energy metabolism along with induction of genes involved in stress responses, such as *sigH*, *sigB*, *groEL2* and *dnaJ*<sup>55</sup>.

Hypoxia is considered to be one of the most important changes for *M.tb* to deal with when it enters the M $\Phi$  and also when it resides within the granuloma during chronic infection. This is reflected by the numerous studies investigating gene regulation in *M.tb* exposed to reduced oxygen tension<sup>95 96 97 98 99 30</sup>. The two-component response regulator *devR-devS* (*dosR-dosS*; *Rv3133c/Rv3132c*) was originally shown to be up regulated during hypoxia along with a regulon of around 50 genes including *hspX* ( $\alpha$ -crystallin), *fdxA*, *narX*, *narK2*, *bfrB* and *ctpF*<sup>95 96</sup>. Later studies confirmed and extended these results using controlled oxygen depletion in a chemostat<sup>97</sup>. In addition to up-regulation of the *devR* regulon, six of the ten contiguous mycobactin synthesis genes (*mbtA-L*) were up-regulated. A comparison of aerobic cultures of *M.tb* with microaerophilic and anaerobic cultures<sup>100</sup> revealed similarities and differences between the two gene sets from the hypoxic *M.tb*<sup>98</sup>. Lysine dehydrogenase, nitrate reductase and  $\alpha$ -crystallin were induced in both treatments, as were genes of the *devR* regulon and fatty acid metabolism genes including *fadD26*. Induction of the fumarate reductase operon (*frdA-D*) following hypoxia has been shown for *M.tb*, presumably as an adaptation to anaerobic respiration<sup>30</sup>. Interestingly, while *M.tb* grown under hypoxic conditions was more virulent in a guinea pig model<sup>97</sup>, a *devR* deletion mutant had no decreased virulence in a murine model of TB<sup>99</sup>, suggesting that the *devR* regulon may be redundant with regards bacterial virulence in mice. Additional analysis showed that induction of the *devR* regulon is transient and is replaced by an “enduring hypoxic response” of 230 genes induced at 4 and 7 days of hypoxia. A meta-analysis of the transcriptional network (origins) differentially expressed in response to hypoxia over time<sup>101</sup>, identified the *devR*, *sigD* and Rv0494 as early origins (response on or before day 6 of hypoxia). Intermediate (day 8-14) origins included *furB/zur*, *crp*, *sigH*, *kstR* and *sigE-mprA*; while *nadR*, Rv1956 and *hrcA* were late origins (>20 days after hypoxia).

#### EXPOSURE TO POTENTIAL MYCOBACTERIACIDAL AGENTS AND PROCESSES

Invading microbes, including pathogens like *M.tb*, can be killed by MMs in a number of ways<sup>102</sup>. M $\Phi$  killing mechanisms that have been shown to be deleterious to *M.tb* include fatty acids<sup>103</sup>, defensins<sup>104</sup> ATP-mediated killing<sup>105</sup>, autophagy<sup>106 107</sup>, glutathione and nitrosogluthathione<sup>108</sup>, p47 GTPases<sup>109</sup> lysosomal ubiquitin derived peptides<sup>110</sup>, the lysosomal enzyme beta-hexosaminidase<sup>111</sup> and the induction of endosomal maturation into phagolysosomes<sup>112 113 114 82 115 86</sup>. However, the major mechanisms that have repeatedly been described as having

antimycobacterial activity are the production of ROI and RNI<sup>116 117 118 119</sup> and the induction of apoptotic cell death of infected MMs<sup>120 121 122 123</sup>.

**ROI:** MΦ produce ROI (superoxide anion, hydrogen peroxide, singlet oxygen) during the ingestion of many types of particles<sup>124 116</sup>. ROI are able to kill mycobacteria<sup>125 126</sup>. In addition, it has been shown that *phox* knockout mice, which are unable to produce ROI, are more susceptible to *M.tb* than their wild type littermates<sup>127 128</sup>, demonstrating the involvement of ROI in killing *M.tb*. Moreover, a *katG* knockout strain of *M.tb* was markedly attenuated in wild-type C57Bl/6 mice and iNOS knockout mice (that can not make RNI), yet was indistinguishable from wild-type *M.tb* in its ability to replicate and persist in *phox* knockout mice<sup>129</sup>. As there is some redundancy in *phox*<sup>116</sup>, it is most probable that ROI play a role in the killing of intracellular mycobacteria but that this is not an exclusive role.

**RNI:** RNI (nitric oxide, the nitrite ion, nitrous anhydride and peroxynitrite) are considered to be involved in the killing of mycobacteria and other microbes by MΦs, either independently or in conjunction with ROI<sup>116 117 130 131</sup>. RNI are produced by MΦs in response to multiple signals and, like ROI, are generally considered to be the result of MΦ activation by a priming (e.g IFNγ) and an activating signal (e.g.LPS )<sup>117</sup>. Studies using iNOS knockout mice<sup>132 127 133</sup> demonstrated that RNI are an important component of the antimycobacterial armature of the murine MΦ though, like ROI, there is some redundancy in iNOS<sup>116 134</sup>. Mice with a combined deficiency in *Phox* and iNOS die of spontaneous microbial infections, illustrating the need for at least one system to be present to fight infection<sup>135</sup>. Recent studies have shown that RNI do not gain access to the mycobacterial phagosome<sup>136</sup> and there is evidence that RNI do not mediate the killing of *M.tb* in MΦs via a direct action on the intracellular bacteria<sup>137</sup>. Thus, the mechanism of RNI toxicity to intracellular *M.tb* is more complex than was first thought. There is some controversy as to whether MΦ production of RNI is a killing mechanism that can be employed by human MΦs<sup>138 139 140</sup>, however, evidence is accumulating that demonstrates that human phagocytes can produce RNI, albeit under tighter regulation than in the murine system<sup>141 142 143 144 145 119</sup>. Thus, whether RNI actually play a role in killing *M.tb* in human MΦs is still unresolved.

**Apoptosis:** Apoptosis is the process of programmed cell death involving a cascade of intracellular signals that results in the death of damaged, infected, or redundant cells without inducing an inflammatory response and the associated damage of surrounding cells. Ligation of apoptotic triggers, such as TNFα, with its receptor, TNFR, leads to the activation of caspases that are responsible for the regulated disassembly of the cell into apoptotic bodies<sup>146</sup>. More recently it has been shown that there are both caspase dependent<sup>147</sup> and independent<sup>148</sup> mechanisms of apoptosis within MΦ that can successfully control intracellular microbes. It has been shown that apoptosis is a mechanism that is employed by the host to eliminate MΦs infected with mycobacteria<sup>149 150 151 152 123</sup>. Moreover mycobacteria (and other pathogens) have been demonstrated to be able to inhibit the apoptotic pathways of MΦs, at least to some extent, resulting in survival of the intracellular bacteria<sup>153 154 152 155</sup>. Recent studies have identified mycobacterial genes (*nuoG* and *secA2*) that are involved in inhibiting apoptosis<sup>156 157</sup>, confirming the importance of apoptosis in MΦ control of *M.tb*.

Changes to the transcriptome of *M.tb* exposed to multiple ROI and RNI donors have been investigated. Hydrogen peroxide was shown to induce the *devR* regulon, but at relatively low levels compared to other stress conditions, including nitrosoglutathione<sup>158</sup>. As would be expected, the catalase gene, *katG* is upregulated following H<sub>2</sub>O<sub>2</sub> treatment<sup>159</sup>. Exposure of *M.tb* to RNI regulated a similar set of genes as did hypoxia, including *devR/devS*, *nrdZ*, *narK2*, *narX*,



*pfkB*, *hspX*, *ctpF* and a number of hypothetical proteins<sup>160 161</sup>. An alternative approach, comparing transcriptomes of *M.tb* in iNOS knockout MΦs versus wild type MΦs gave similar results and identified an NO-induced response that represented a shift from aerobic to anaerobic respiration (induction of the *devR* regulon) and increased iron scavenging and utilisation of fatty acids as a carbon source.<sup>56</sup> Unfortunately, no information is available for gene regulation of *M.tb* within apoptotic MΦ, although, as both ROI and RNI induce apoptosis, the studies comparing *M.tb* gene expression in the presence of these toxic molecules may reflect what would be found for bacteria within apoptotic MΦ.

#### THE INDUCTION OF ADAPTIVE IMMUNITY AND THE MOVE TO A DORMANCY PROFILE.

The early interactions of *M.tb* with MΦ are generally considered to take place in the absence of any adaptive immune response. Rather, the bacteria are exposed to elements of innate immunity, of which MΦ are an important component. Although innate immunity exerts a level of control on the replication of *M.tb* during the early acute phase of disease<sup>123 162</sup>, it is only when antigen specific adaptive immunity is initiated that bacterial replication dramatically slows down and *M.tb* enters into a chronic stage of disease. This is characterised by minimal increase, or even slow decline, in the bacterial burden and a move towards a physiological state that is often referred to as dormancy. That this control of bacterial replication is mediated by the immune system has been demonstrated by studies in knockout mice lacking CD4+ T cells, or with defects in the IFN $\gamma$  activation pathway (IFN $\gamma$  production, IFN $\gamma$  receptor, IL-12), or with defects in TNF $\alpha$  or RNI production. These mice are significantly more susceptible to *M.tb* than intact littermates. Other components of the adaptive immune response such as CD8+ T cells are also important<sup>163 164</sup> but do not appear to be as critical.<sup>165 166</sup> From these, and other studies, it is generally accepted that, while there are a number of contributory factors to adaptive immunity to *M.tb*, the most important elements are the production of CD4+ve TH<sub>1</sub> cells that produce IFN $\gamma$ , CD8+ve cytolytic T lymphocytes (also IFN $\gamma$  producers) and TNF $\alpha$ <sup>167 168 169</sup>. The most important way these critical elements mediate control of bacterial survival and replication is likely through the action of IFN $\gamma$  and TNF $\alpha$  on the MΦ and the granuloma. Both IFN $\gamma$  and TNF $\alpha$  can act alone or in synergy to activate infected MΦs, resulting in an increase in phagosomal maturation towards a phagolysosome<sup>170</sup>, increased ROI and RNI production by the MΦ<sup>116</sup> and increased apoptosis<sup>171 172-174</sup>, ultimately resulting in a MΦ intracellular environment that is non-permissive for mycobacterial replication. To counter this response by the host, *M.tb* can down-regulate the induction of cytokines involved in the activation of MΦ. Studies using purified mycobacterial components or directed cell wall mutants of *M.tb* indicate that survival of intracellular *M.tb* is linked to the ability to down-regulate pro-inflammatory cytokines such as IL-12, IL-6 and TNF $\alpha$ <sup>172-174</sup> and inhibit the activation of MΦ by IFN $\gamma$ <sup>175</sup>. However, even though *M.tb* can modulate the cytokine response of the host, it is evident from animal infection models that this is not sufficient to prevent the activation of MΦ to a state capable of controlling *M.tb* growth and survival *in vivo*.

In response to IFN $\gamma$ -mediated activation of the MΦ, *M.tb* alters its physiology to enter into a dormancy phase characterised by a drastically reduced level of replication<sup>176</sup> or a balance between slow replication and killing<sup>162</sup>, use of lipids as a carbon source through the glyoxalate shunt<sup>177</sup> and a shift towards microaerophilic respiration<sup>178</sup>. The IFN $\gamma$ -mediated maturation of the phagosome will also lead to further nutrient deprivation and hypoxia. These signals are thought to lead to a dormancy regulon that enables *M.tb* to persist inside the MΦ and the

granuloma. Meta-analysis of microarray studies modelling dormancy<sup>179</sup>, indicated several trends including up-regulation of 81% of the *devR* regulon, down-regulation of over 50% of the 30S and 50S ribosomal-protein genes and down regulation of ATP synthesis. This is indicative of a bacterium slowing down protein synthesis and shifting towards alternative electron acceptors such as nitrate or fumarate during respiration. *M.tb* is nominally an aerobe but is able to switch to a microaerophilic/anaerobic mode of respiration as indicated by the up-regulation of *fdxA*, *narX*, *narK2* and the *frd* operon. Other features of the dormancy regulon include the further induction of genes involved in the utilization of fatty acids as a carbon source and genes involved in mycolic acid modifications, indicating modification of the cell envelope<sup>56</sup>. As RNI are only produced by activated MΦ, it has been suggested that they may be an important trigger of dormancy in *M.tb* following the onset of adaptive immunity. RNI can inhibit respiration in bacteria and it is therefore not surprising that the RNI induced dormancy regulon closely resembled that induced by hypoxia and nutrient starvation as represented by the *devR* regulon<sup>160</sup>  
161 180.

## CONCLUSIONS AND CAVEATS

The move of *M.tb* from extracellular to intracellular residence within the MΦ is a major change in environment, requiring major adjustments to enable the bacterium to survive and flourish. We have reviewed those environmental changes and the accompanying bacterial adaptations necessary for *M.tb* to succeed as a pathogen at the level of gene expression. While there is no complete agreement among the numerous studies as to the identity of genes regulated by this adaptation, nor for that matter as to the identity of the elements of the MΦ environment that induce this regulation or indeed the multiple microenvironments that may be encountered by infecting bacilli, there is a level of consensus. It is generally agreed that the move to an intramacrophage location will result in *M.tb* having to deal with changes in oxygen tension, nutrient depletion, increased acidity and exposure to immune defences such as ROI, RNI, apoptosis and phagosomal maturation. During the early, acute stage of infection, *M.tb* bacilli are likely exposed to the innate immune response which includes such elements as complement, NK cells, defensins, neutrophils and resident MΦs. Although these clearly have an effect, some bacteria seem able to adapt to or avoid these challenges and replicate relatively successfully. Following the development of adaptive immunity, the infection shifts into a chronic stage where the host's defences are more effective at controlling bacterial replication and survival. The bacteria are walled off in a granuloma and go into a dormant or latent phase that is maintained predominately by host IFNγ and TNFα.

*M.tb* responds to these changes by rapidly changing its transcriptome. Intracellular residence, hypoxia, nitrites and starvation all seem to induce genes characterized by the dormancy regulon that enable bacilli to undergo anaerobic respiration, utilize lipids as a carbon source and modify the cell envelope. Many studies have shown that the *devR* (*dosR*) regulon is heavily represented in all treatments that lead to a dormancy profile in *M.tb*, suggesting that these genes are very important in the continued survival of intracellular bacteria. However, recent data suggests that the *devR* regulon is not necessary for bacterial virulence and appears to be transiently expressed during early hypoxia (< 24 hours) and that an enduring hypoxic response of 230 genes contains key factors needed for *M.tb* persistence<sup>99</sup>. The *hspX* gene encoding the 16kD α-crystallin chaperonin is also commonly up-regulated in *M.tb* on entering MΦs and moving into dormancy. Indeed, its expression is often used as an indication of a dormant state. However, it has been shown that letting cultures stand for 30 minutes can result in 100 fold increases in *hspX* expression<sup>158</sup>, which suggests that dormancy is not the only trigger leading to increased production of this chaperonin. It appears that *hspX*, along with many other genes differentially expressed intracellularly, are inextricably linked to a slowing of *M.tb* replication after multiple stresses. Many of the studies investigating gene expression by *M.tb* in MΦs have modelled the intracellular environment in axenic culture in order to determine the effect of a single variable over time, and to avoid the complications of isolating RNA from intracellular bacteria. This has often led to exposing the bacteria to a single trigger hypothesised to reflect the phagosomal

environment, such as acidity, hypoxia, H<sub>2</sub>O<sub>2</sub> or nitrite. While this reductionist approach has produced valuable information, it has to be remembered that the intracellular *M.tb* bacilli are not exposed to each of these conditions in isolation and it is perhaps better to try and determine the bacteria's response to intracellular life by actually looking at bacteria within the MΦ. Perhaps more informative are studies aimed at identifying the functional significance of isolated elements of the host response by using MΦ with single gene deletants (for example, ROI or RNI) to yield single pathway inferences. Studies designed to investigate the multi-factorial nature of *M.tb* infection should provide new perspectives, but will have to negotiate increasingly complex networks of interaction.

The distinction between intracellular microenvironments permissive and non-permissive for *M.tb* growth has also been modelled by comparing the intracellular transcriptomes of different mycobacterial species and in different phagocytic cells, for example *M.tb* compared to BCG<sup>55</sup>, and *M.tb* infection of MΦ compared to dendritic cells<sup>60</sup>. This raises an important element in studying the transcriptome of *M.tb* within the intracellular environment. The actual model utilised i.e. the mycobacterial species and phagocyte used will have significant impact on the results obtained. Many studies have been performed using *M. bovis* BCG as a model for *M.tb* because of the high degree of homology between the two bacterial genomes, coupled with the relative ease with which BCG can be used experimentally (it does not require containment level 3 laboratories). However, BCG lacks an important virulence region called RD1<sup>181</sup>, is attenuated in humans and does not result in the same pathogenesis as *M.tb*. While the genome sequence of *M.tb* and *M. bovis* are very similar, the transcriptomes of the 2 species show several differences<sup>182</sup> demonstrating the need to use *M.tb* in expression analysis of *M.tb* in MΦ. The other major source of variation in trying to understand the intracellular phenotype of *M.tb* is the choice of MΦ infection model itself. Commonly used models are the murine bone marrow-derived MΦ (BMM)<sup>56 58 55</sup>, human monocyte-derived MΦ (MDM)<sup>57 60</sup> and the human MΦ-like cell line THP-1<sup>59</sup>. Each of these models has strengths and weaknesses. For instance, when using MDM, significant variation between the MΦs from individual blood donors means that greater numbers of samples are required before variation is reduced enough to characterize specific patterns. The use of THP-1 cells overcomes this variation but introduces problems associated with it being a cell line that is not truly representative of a MΦ. Perhaps the BMM is the best model as it represents a reasonably uniform population of MΦ that are derived from *ex vivo* cells and, unlike human MΦ models, results obtained in BMM *in vitro* can be further tested in infection models *in vivo*. However, a long-standing issue with the use of murine MΦs is the strong inducible RNI response seen following immune activation. This has been identified as one of the major determinants resulting in the induction of the *M.tb* intracellular dormancy program. While it is becoming increasingly clear that human MΦs can make RNI<sup>119</sup>, there are significant differences between human and mouse MΦ RNI production. Additionally, strain variation between mice may also affect the results using BMM.

Finally, it is necessary to ensure that comparisons between data sets of genes take into consideration the behaviour of *M.tb* in the MΦ model used. For instance, *M.tb* usually replicates efficiently in untreated "resting" murine BMM over 7 days *in vitro* and IFNγ treatment can slow this replication rate but does not result in bacterial elimination<sup>183 109 123</sup>. In one of the seminal studies on *M.tb* gene expression in MΦ<sup>56</sup>, no replication was seen in resting murine BMM MΦ, and IFNγ stimulation resulted in bacterial elimination. While there are many possible explanations for this disparity, it is important to ensure that variations in the models used are considered when comparing *M.tb* intracellular RNA profiles. An additional source of variation, particularly applicable to the use of microarray platforms to study changes in gene expression, is the choice of comparator conditions. Typically, intracellular transcriptional profiles are compared to log phase axenic bacilli, or bacilli resuspended in tissue culture media alone. Transcriptional differences between these bacterial populations will necessarily affect the interpretation of intracellular RNA profiles, and make the meaningful comparison of intracellular datasets problematic. Figure 3 displays one such comparison of four publicly available microarray

1 datasets describing the global response of *M.tb* to macrophage infection. Significant technical  
2 variation between studies makes the direct comparison of transcriptional profiles unwise;  
3 however the similarities between models are clearly visible. Interestingly, it is also possible to  
4 identify host-specific patterns that are represented across several datasets. Further work will be  
5 required to ascertain the significance of such findings.

#### 6 7 8 FUTURE PERSPECTIVES

9 An understanding of the transcriptome of *M.tb* in the MΦ is developing and has identified a  
10 number of physiological changes the bacterium must undergo to survive within its specialised  
11 phagosome. Most of this information comes from use of *in vitro* models of *M.tb* in MΦs or of  
12 axenic bacteria exposed to conditions considered to reflect the intracellular environment. There  
13 have been some important studies investigating gene expression of *M.tb* in whole animal models  
14 <sup>184 36 185 186 187 39</sup> and humans <sup>188 38 189</sup>. While *in vitro* studies are informative and an essential  
15 step in understanding the physiology of intracellular *M.tb*; the complexity of the environment in  
16 the host with the associated myriad of potential triggers of bacterial gene regulation mean that a  
17 full understanding of the intracellular transcriptomes of *M.tb* during its ongoing pathogenesis and  
18 continued survival in multiple microenvironments in the host will only be fully understood  
19 through *in vivo* investigations. Gene expression profiling of *M.tb* bacilli extracted from distinct  
20 stages of infection using human tissue and animal models will be required. Although the mouse,  
21 guinea pig and rabbit are useful models of tuberculosis, the primate model of tuberculosis better  
22 reflects disease in humans <sup>190 191</sup>. Although challenging, studying the transcriptome of *M.tb*  
23 during the development of tuberculosis in the primate may give a greater understanding of the  
24 disease process in humans.

25  
26 There are still areas of the host/pathogen interaction in tuberculosis that will greatly benefit  
27 from *in vitro* studies. One area that has not been studied is the gene expression of *M.tb* exposed  
28 to apoptotic MΦs. Apoptosis is a major killing mechanism of MΦs that has been shown to be  
29 very important in the control of *M.tb* <sup>149 123 192</sup>. *M.tb* has evolved mechanisms to inhibit apoptosis  
30 (which is protective for the host); instead *M.tb* induces necrosis in infected MΦ, which is  
31 beneficial to bacterial survival <sup>153 157 156 193 123</sup>. However, although gene expression by MΦs  
32 undergoing apoptosis or necrosis following infection with *M.tb* has been investigated <sup>194</sup>, no  
33 study of the bacterial transcriptome in apoptotic versus necrotic MΦs has been undertaken. As  
34 inhibition of apoptosis is a critical survival mechanism for *M.tb*, it will be of great interest to  
35 define the transcriptional basis of this process.

36  
37 Another area that has not been extensively investigated at the level of bacterial gene  
38 expression is the comparison of virulent and avirulent *M.tb* in MΦs. Some work has been done  
39 comparing *M.tb* with the attenuated vaccine strain, *M.bovis* BCG <sup>55</sup> but there are significant  
40 differences between *M.tb* and BCG which makes comparing the two strains difficult. Recent  
41 work has compared the transcriptomes of a virulent strain of *M.tb* (H37Rv) with an avirulent  
42 strain (H37Ra) during growth in broth and identified a number of genes that are differentially  
43 regulated in the attenuated H37Ra that include members of the dormancy regulon and a number  
44 of genes connected to *phoP* expression <sup>195 196 197</sup>. Whether these differences are repeated when  
45 H37Rv and H37Ra are compared when inside MΦs awaits investigation, although one recent  
46 study has identified *frdA-D* among differentially regulated genes between intracellular H37Rv  
47 and H37Ra <sup>30</sup>. Transcriptional profiling of *M.tb* knockout mutants with specific *in vivo*  
48 phenotypes of compromised virulence <sup>198</sup> may become an important tool for defining the  
49 functional significance of single gene products and understanding the molecular determinants  
50 defining *M.tb* pathogenicity and virulence.

51  
52 The intracellular transcriptomes of *M.tb* may help us to understand the physiological and  
53 metabolic states required by *M.tb* to successfully complete a cycle of human infection. They  
54 may also be utilised as a bioprobe to discover and define microenvironments encountered by

1 bacilli during infection. Additionally, they may facilitate the discovery of virulence mechanisms  
2 that enable *M.tb* bacilli to modulate the host immune response. The identification of genes  
3 induced intracellularly suggests that they are important for the survival of *M.tb* in the host  
4 environment, and would therefore be possible candidate targets for therapeutic discovery<sup>179</sup>. As  
5 novel therapeutics are critically needed to control tuberculosis, especially with XDR *M.tb*  
6 becoming so prevalent<sup>2</sup>, any strategy that facilitates drug discovery should be actively pursued.  
7 Additionally, these targets could act as potential candidates in rational vaccine design either to  
8 replace or augment the current BCG vaccine.

## 11 EXECUTIVE SUMMARY

### 12 Overview of the infectious process

- 13 ■ *M.tb* binds to MΦ receptors and is ingested into a phagosome.
- 14 ■ *M.tb* inhibits the development of the phagosome, holding it at the early endosome stage.
- 15 ■ *M.tb* induces the production of cytokines that result in a TH<sub>1</sub> adaptive immune response.
- 16 ■ Through the activity of IFNγ and TNFα, infected MΦ become activated and are better  
17 able to control the replication of *M.tb* through the action of Reactive Nitrogen  
18 Intermediates, Reactive Oxygen Intermediates and apoptosis.
- 19 ■ *M.tb* then moves to a dormant state, often within a granuloma, and changes its physiology  
20 to combat the variation in oxygen tension, low carbohydrates and MΦ killing  
21 mechanisms.

### 23 Binding and uptake (movement from extracellular to intracellular)

- 24 ■ Non-opsonic uptake of *M.tb* is important in the serum free environment of the lung.
- 25 ■ *M.tb* moderates its non-opsonic uptake by MΦ.
- 26 ■ Transcriptional responses of *M.tb* begin within minutes of ingestion and include members  
27 of the *devRS* regulon and the *phoPR* regulon
- 28 ■ A comparison of numerous microarray experiments identified common functionalities in  
29 *M.tb* regulated gene sets indicating a move to an environment low in carbohydrate, rich in  
30 lipid, low in oxygen and with toxic molecules present.

### 32 Residence in a phagosome and the associated low nutrients, increased acidity and hypoxia.

- 33 ■ *M.tb* maintains the phagosome at a mildly acidic pH and with markers indicating arrest at  
34 the early endosome stage with access to extracellular nutrients.
- 35 ■ Nutrient deprivation, exposure to acidic pH and hypoxia have been used to model the  
36 phagosome.
- 37 ■ This resulted in *M.tb* regulation of genes involved in oxidation of fatty acids, aerobic  
38 respiration, energy metabolism and lipid biosynthesis.
- 39 ■ The *devRS* and *phoPR* regulons were upregulated.

### 41 Exposure to potential mycobacteriacidal agents and processes

- 42 ■ The major mycobacteriacidal mechanisms appear to be RNI, ROI and apoptosis.
- 43 ■ ROI induces the upregulation of *katG*.
- 44 ■ RNI induce the *devRS* regulon.

### 46 The induction of adaptive immunity and the move to a dormancy profile.

- 47 ■ Adaptive immunity is essential for the optimal control of *M.tb*.
- 48 ■ The most important components of the effective adaptive response are IFNγ producing  
49 CD4+ve TH<sub>1</sub> cells, CD8+ve cytolytic T cells and TNFα.
- 50 ■ IFNγ activates MΦ to better control the survival and replication of *M.tb*.
- 51 ■ This induces dormancy in *M.tb*, characterised by lower rates of replication, use of lipids  
52 as a carbon source and a shift to microaerophilic respiration.
- 53 ■ The dormancy regulon is dominated by the *devRS* regulon, genes involved in mycolic  
54 acid modifications and genes involved in the utilization of fatty acids.

## ACKNOWLEDGEMENTS

We would like to thank Lisa Thorson for preparing Figure 1. RWS is the recipient of a Child & Family Research Institute Investigatorship Award. SJW is currently funded by a European Union 6th Framework Programme 'New Medicines for Tuberculosis', NM4TB (project number LSHP-CT-2005-018923).

## Figure Legends

### **Figure 1. Interaction of *M.tb* with macrophages.**

*M. tuberculosis* (*M. tb*) expressing a variety of Pathogen Associated Molecular Patterns (PAMPs), is recognized by Pattern Recognition Receptors (PRR) on host macrophages leading to BINDING and then UPTAKE. During binding and uptake Reactive Oxygen Intermediates (ROI), Reactive Nitrogen Intermediates (RNI) and Cytokines are released by the macrophage. The INNATE immune response drives the ADAPTIVE response and, depending on the cytokine profile induced, will result in TH1 or TH2 and pro- or anti- inflammatory responses. The bacteria reside within an EARLY PHAGOSOME in which Rab5, coronin 1 and proCathepsin D (PCD) are found. The mycobacterial phagosome also has a relatively high pH and has access to the extracellular milieu as shown by the cycling of transferrin. Mycobacterial antigens are processed in some, as yet, poorly defined way leading to PRESENTATION. Under some circumstances, mycobacteria are killed (xxx) and are then found in LATE PHAGOSOMES which contain Lamp 1, Rab 7 and cathepsin D and also have a V-ATPase proton pump which acidifies the phagosome. The killing of intracellular mycobacteria can be mediated by RNI, ROI or through the INDUCTION OF APOPTOSIS which can be triggered by ROI, RNI and cytokines such as TNF $\alpha$ .

### **Figure 2. Dissecting the *M.tb* intracellular transcriptional profile.**

535 genes identified as significantly differentially expressed in two or more studies describing the *M.tb* response to macrophage infection (A-D). Genes identified (by the study authors) to be significantly induced (marked in red) or repressed (marked in blue) from *in vitro* axenic models broadly representing low pH (E-H), hypoxia (I-M), nutrient starvation (N-P) and oxidative stress (Q-S) are mapped alongside the intracellular response. Conditions are detailed as columns, genes as rows. The asterisked column represents an additive model describing the sum induction or repression of each gene from the axenic conditions. This rudimentary model shows the extent to which current *in vitro* modelling helps to define the *M.tb* intracellular profile. For example, Box 1 highlights the hypoxic response as an integral element of the *M.tb* intracellular response. Box 2 marks a cluster of genes differentially regulated by multiple *in vitro* environments. Box 3 shows the repression of ATP synthase and NAD dehydrogenase I (*atp* and *nuo*) gene families intracellularly. All *M.tb* axenic conditions relative to log phase untreated controls unless otherwise stated. (A) Murine bone marrow-derived M $\Phi$  (BMDM) <sup>56</sup>. (B) Human monocyte-derived M $\Phi$  (MDM), <sup>60</sup>. (C) THP-1 human M $\Phi$ -like cells (THP-1), <sup>92</sup>. (D) Murine BMDM relative to media control, <sup>55</sup>. (E) Acid shock, <sup>182</sup>. (F) Low pH, <sup>93</sup> (G) pH 6.5, <sup>55</sup> (H) pH 5.5 <sup>55</sup>. (I) Hypoxia, <sup>97</sup> (J) Non-replicating persistence-1 <sup>189</sup> (K) Non-replicating persistence-2 <sup>189</sup> (L) Non-

replicating persistence or stationary phase<sup>180</sup> (M) Enduring hypoxic response<sup>99</sup> (N) Low glucose<sup>88</sup> (O) Nutrient starvation<sup>89</sup> (P) Low copper<sup>199</sup> (Q) Reactive nitrogen intermediates<sup>160</sup> (R) Oxidative stress<sup>200</sup> (S) SDS stress<sup>201</sup>

### Figure 3 The *M.tb* response to the intracellular environment.

The transcriptional profile of 1121 genes (>2 fold differentially expressed in >2 conditions) were extracted, combined and clustered from four publicly available microarray datasets. Conditions/timepoints are displayed as columns, genes as rows. Red colouring indicates induction, blue repression compared to aerobic log phase growth (unless otherwise stated). MDM, human monocyte-derived MΦ infection model<sup>60</sup>. BMDM, naïve and IFNγ-activated murine bone marrow-derived MΦs<sup>56</sup>. THP, THP-1 human MΦ-like cell line infection model<sup>92</sup>. BM, murine bone marrow-derived MΦs infected with *M.tb* relative to bacilli resuspended in tissue culture media<sup>55</sup>. Box A – genes differentially expressed after infection of human-derived cells only (MDM/THP). Box B – temporal response to infection observed in murine-derived cells only (BMDM/BM).

### REFERENCES

1. Corbett EL, Watt CJ, Walker N, et al. The growing burden of tuberculosis: Global trends and interactions with the HIV epidemic. *Arch Intern Med* 2003;**163**(9):1009-1021.
2. Chan ED, Iseman MD. Multidrug-resistant and extensively drug-resistant tuberculosis: a review. *Curr Opin Infect Dis*. 2008;**21**(6):587-595.
3. Malik S, Schurr E. Genetic susceptibility to tuberculosis. *Clin Chem Lab Med* 2002;**40**(9):863-868.
4. Chackerian AA, Behar SM. Susceptibility to *Mycobacterium tuberculosis*: lessons from inbred strains of mice. *Tuberculosis* 2003;**83**(5):279-285.
5. Doffinger R, Patel SY, Kumararatne DS. Host genetic factors and mycobacterial infections: lessons from single gene disorders affecting innate and adaptive immunity. *Microbes Infect* 2006;**8**(4):1141-1150.
6. Kramnik I. Genetic dissection of host resistance to *Mycobacterium tuberculosis*: the sst1 locus and the Ipr1 gene. *Curr Top Microbiol Immunol*. 2008;**321**:123-48.
7. Collins FM, Smith MM. A comparative study of the virulence of *Mycobacterium tuberculosis* measured in mice and guinea pigs. *Am Rev Respir Dis* 1969;**100**:631-639.
8. Dunn PL, North RJ. Virulence ranking of some *Mycobacterium tuberculosis* and *Mycobacterium bovis* strains according to their ability to multiply in the lungs, induce lung pathology, and cause mortality in mice. *Infect Immun* 1995;**63**(9):3428-3437.
9. Guilhot C, Otal I, Vanrompaey I, Martin C, Gicquel B. Efficient transposition in mycobacteria - construction of *Mycobacterium smegmatis* insertional mutant libraries. *J Bacteriol* 1994;**176**(2):535-539.
10. Pelicic V, Jackson M, Reyrat JM, Jacobs WR, Gicquel B, Guilhot C. Efficient allelic exchange and transposon mutagenesis in *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U S A* 1997;**94**(20):10955-10960.
11. Cox JS, Chen B, McNeil M, Jacobs WR. Complex lipid determines tissue specific replication of *Mycobacterium tuberculosis* in mice. *Nature* 1999;**402**(6757):79-83.
12. Camacho LR, Ensergueix D, Perez E, Gicquel B, Guilhot C. Identification of a virulence gene cluster of *Mycobacterium tuberculosis* by signature-tagged transposon mutagenesis. *Mol Microbiol* 1999;**34**(2):257-267.

- 1 \*13. Sassetti CM, Rubin EJ. Genetic requirements for mycobacterial survival during infection.  
2 *Proc Natl Acad Sci U S A* 2003;**100**(22):12989-12994.
- 3 \*14. Rengarajan J, Bloom BR, Rubin EJ. Genome-wide requirements for *Mycobacterium*  
4 *tuberculosis* adaptation and survival in macrophages. *Proc Natl Acad Sci U S A*  
5 2005;**102**(23):8327-8332.
- 6 These two references identified essential genes necessary for the survival of *Mycobacterium*  
7 *tuberculosis* *in vivo* within mice and intracellularly within macrophages.  
8
- 9 15. Rindi L, Lari N, Garzelli CA. Search for genes potentially involved in *Mycobacterium*  
10 *tuberculosis* virulence by mRNA differential display. *Biochem Biophys Res Commun*  
11 1999;**258**(1):94-101.
- 12 16. Plum G, Clark-Curtiss JE. Induction of *Mycobacterium avium* gene expression following  
13 phagocytosis by human macrophages. *Infect Immun* 1994;**62**(2):476-483.
- 14 17. Plum G, Brenden M, Clark-Curtiss JE, Pulverer G. Cloning, sequencing, and expression of  
15 the mig gene of *Mycobacterium avium*, which codes for a secreted macrophage-induced  
16 protein. *Infect Immun* 1997;**65**(11):4548-4557.
- 17 18. Graham JE, Clark-Curtiss JE. Identification of *Mycobacterium tuberculosis* RNAs  
18 synthesized in response to phagocytosis by human macrophages by selective capture of  
19 transcribed sequences (SCOTS). *Proc Natl Acad Sci U S A* 1999;**96**(20):11554-11559.
- 20 19. Dhandayuthapani S, Via LE, Thomas CA, Horowitz PM, Deretic D, Deretic V. Green  
21 fluorescent protein as a marker for gene expression and cell biology of mycobacterial  
22 interactions with macrophages. *Mol Microbiol* 1995;**17**(5):901-912.
- 23 20. Barker LP, Brooks DM, Small PLC. The identification of *Mycobacterium marinum* genes  
24 differentially expressed in macrophage phagosomes using promoter fusions to green  
25 fluorescent protein. *Mol Microbiol* 1998;**29**(5):1167-1177.
- 26 21. Triccas JA, Berthet FX, Pelicic V, Gicquel B. Use of fluorescence induction and sucrose  
27 counterselection to identify *Mycobacterium tuberculosis* genes expressed within host  
28 cells. *Microbiology* 1999;**145**(10):2923-2930.
- 29 22. Butcher PD. Microarrays for *Mycobacterium tuberculosis*. *Tuberculosis* 2004;**84**(3-4):131-  
30 137.
- 31 23. Talaat AM, Hunter P, Johnston SA. Genome-directed primers for selective labeling of  
32 bacterial transcripts for DNA microarray analysis. *Nat Biotechnol* 2000;**18**(6):679-682.
- 33 24. Waddell SJ, Laing K, Senner C, Butcher PD. Microarray analysis of defined *Mycobacterium*  
34 *tuberculosis* populations using RNA amplification strategies. *BMC Genomics*. 2008;**9**:94.
- 35 25. Brosch R, Gordon SV, Billault A, et al. Use of a *Mycobacterium tuberculosis* H37Rv  
36 bacterial artificial chromosome library for genome mapping, sequencing, and comparative  
37 genomics. *Infect Immun* 1998;**66**(5):2221-2229.
- 38 26. Brosch R, Philipp WJ, Stavropoulos E, Colston MJ, Cole ST, Gordon SV. Genomic analysis  
39 reveals variation between *Mycobacterium tuberculosis* H37Rv and the attenuated *M.*  
40 *tuberculosis* H37Ra strain. *Infect Immun* 1999;**67**(11):5768-5774.
- 41 27. Gordon SV, Brosch R, Billault A, Garnier T, Eiglmeier K, Cole ST. Identification of variable  
42 regions in the genomes of tubercle bacilli using bacterial artificial chromosome arrays.  
43 *Mol Microbiol* 1999;**32**(3):643-655.
- 44 28. Brosch R, Gordon SV, Eiglmeier K, Garnier T, Cole ST. Comparative genomics of the  
45 leprosy and tubercle bacilli. *Res Microbiol* 2000;**151**(2):135-142.



29. Li AH, Lam WL, Stokes RW. Bacterial artificial chromosome fingerprint arrays for the differentiation of transcriptomic differences in mycobacteria. *J Microbiol Meth* 2008;**75**:416-424.
  30. Li AH, Lam WL, Stokes RW. Characterization of genes differentially expressed within macrophages by virulent and attenuated *Mycobacterium tuberculosis* identifies candidate genes involved in intracellular growth. *Microbiology*. 2008;**154**(Pt 8):2291-2303.
  31. Denis M. Human neutrophils, activated with cytokines or not, do not kill virulent *Mycobacterium tuberculosis*. *J Infect Dis* 1991;**163**:919-920.
  32. Eruslanov EB, Lyadova IV, Kondratieva TK, et al. Neutrophil Responses to Mycobacterium tuberculosis Infection in Genetically Susceptible and Resistant Mice. *Infect. Immun*. 2005;**73**(3):1744-1753.
  33. Bermudez LE, Goodman J. *Mycobacterium tuberculosis* invades and replicates within type II alveolar cells. *Infect Immun* 1996;**64**(4):1400-1406.
  34. Teitelbaum R, Schubert W, Gunther L, et al. The Mcell as a portal of entry to the lung for the bacterial pathogen *Mycobacterium tuberculosis*. *Immunity* 1999;**10**(6):641-650.
  35. Li MS, Waddell SJ, Monahan IM, et al. Increased transcription of a potential sigma factor regulatory gene Rv1364c in *Mycobacterium bovis* BCG while residing in macrophages indicates use of alternative promoters. *FEMS Microbiol Lett* 2004;**233**(2):333-339.
  - \*36. Talaat AM, Lyons R, Howard ST, Johnston SA. The temporal expression profile of *Mycobacterium tuberculosis* infection in mice. *Proc Natl Acad Sci U S A* 2004;**101**(13):4602-4607.
- This reference identifies gene sets that are expressed at different stages of an *in vivo* infection.
37. Karakousis PC, Yoshimatsu T, Lamichhane G, et al. Dormancy Phenotype Displayed by Extracellular *Mycobacterium tuberculosis* within Artificial Granulomas in Mice. *J. Exp. Med*. 2004;**200**(5):647-657.
  - \*38. Rachman H, Strong M, Ulrichs T, et al. Unique transcriptome signature of *Mycobacterium tuberculosis* in pulmonary Tuberculosis. *Infect. Immun*. 2006;**74**(2):1233-1242.
- This reference is the first expression analysis of *Mycobacterium tuberculosis* from clinical lung samples.
39. Kesavan AK, Brooks M, Tufariello J, Chan J, Manabe YC. Tuberculosis genes expressed during persistence and reactivation in the resistant rabbit model. *Tuberculosis* 2009;**89**(1):17-21.
  40. Srivastava V, Jain A, Srivastava BS, Srivastava R. Selection of genes of *Mycobacterium tuberculosis* upregulated during residence in lungs of infected mice. *Tuberculosis* 2008;**88**(3):171-177.
  41. Stokes RW, Haidl ID, Jefferies WA, Speert DP. Mycobacteria-macrophage interactions. Macrophage phenotype determines the nonopsonic binding of *Mycobacterium tuberculosis* to murine macrophages. *J Immunol* 1993;**151**(12):7067-7076.
  42. Stokes RW, Thorson LM, Speert DP. Nonopsonic and opsonic association of *Mycobacterium tuberculosis* with resident alveolar macrophages is inefficient. *J Immunol* 1998;**160**(11):5514-5521.
  43. Stokes RW, Norris-Jones R, Brooks DE, Beveridge TJ, Doxsee D, Thorson LM. The glycan rich outer layer of the cell wall envelope of *Mycobacterium tuberculosis* acts as an anti-phagocytic capsule limiting the association of the bacterium with macrophages. *Infect Immun* 2004;**72**(10):5676-5686.

44. Swartz RP, Naai D, Vogel C-W, Yeager H. Differences in uptake of mycobacteria by human monocytes: a role for complement. *Infect Immun* 1988;**56**(9):2223-2227.
45. Schlesinger LS. Entry of *Mycobacterium tuberculosis* into mononuclear phagocytes. *Curr Top Microbiol Immunol*. 1996;**215**:71-96.
46. Reynolds HY, Newball HN. Analysis of proteins and respiratory cells obtained from human lungs by bronchial lavage. *J Lab Clin Med* 1974;**84**(4):559-573.
47. Mitchell TG, Perfect J. Cryptococcosis in the era of AIDS - 100 years after the discovery of *Cryptococcus neoformans*. *Clin Microbiol Rev*. 1995;**8**(4):515-548.
48. Berger M, Norvell T, Tosi M, Konstan M, Schreiber J. Lack of complement receptor expression by alveolar macrophages (AM) correlates with poor enhancement of phagocytosis of *P. aeruginosa* by complement (C). *Pediatr Res* 1990;**27**(4, part 2):154A.
49. Schlesinger LS, Hull SR, Kaufman TM. Binding of the terminal mannosyl units of lipoarabinomannan from a virulent strain of *Mycobacterium tuberculosis* to human macrophages. *J Immunol* 1994;**152**(8):4070-4079.
50. Peterson PK, Gekker G, Hu S, et al. CD14 receptor-mediated uptake of nonopsonised *Mycobacterium tuberculosis* by human microglia. *Infect Immun* 1995;**63**(4):1598-1602.
51. Gaynor CD, McCormack FX, Voelker DR, McGowan SE, Schlesinger LS. Pulmonary surfactant protein a mediates enhanced phagocytosis of *Mycobacterium tuberculosis* by a direct interaction with human macrophages. *J Immunol* 1995;**155**(11):5343-5351.
52. Melo MD, Catchpole IR, Haggart G, Stokes RW. Utilization of CD11b knockout mice to characterize the role of complement receptor 3 (CR3, CD11b/CD18) in the growth of *Mycobacterium tuberculosis* in macrophages. *Cell Immunol*. 2000;**205**(1):13-23.
53. Fratazzi C, Manjunath N, Arbeit RD, et al. A macrophage invasion mechanism for mycobacteria implicating the extracellular domain of CD43. *J Exp Med* 2000;**192**(2):183-192.
54. Randhawa AK, Ziltener HJ, Merzaban JS, Stokes RW. CD43 is required for optimal growth inhibition of *Mycobacterium tuberculosis* in macrophages and in mice. *J Immunol* 2005;**175**(3):1805-1812.
- \*\*55. Rohde KH, Abramovitch RB, Russell DG. *Mycobacterium tuberculosis* invasion of macrophages: linking bacterial gene expression to environmental cues. *Cell Host Microb* 2007;**2**(5):352-364.
- This reference identifies a profile of *Mycobacterium tuberculosis* infection-linked gene expression over the initial 2 hours of interaction with macrophages.
- \*\*56. Schnappinger D, Ehrt S, Voskuil MI, et al. Transcriptional adaptation of *Mycobacterium tuberculosis* within macrophages: Insights into the phagosomal environment. *J. Exp. Med.*, 2003: 693-704.
- This was a seminal publication investigating the transcriptome of *Mycobacterium tuberculosis* within macrophages.
57. Haydel SE, Clark-Curtiss JE. Global expression analysis of two-component system regulator genes during *Mycobacterium tuberculosis* growth in human macrophages. *FEMS Microbiol Lett* 2004;**236**(2):341-347.
58. Rachman H, Strong M, Schaible U, et al. *Mycobacterium tuberculosis* gene expression profiling within the context of protein networks. *Microb Infect* 2006;**8**(3):747-757.

59. Fontan P, Aris V, Ghanny S, Soteropoulos P, Smith I. The global transcriptional profile of *Mycobacterium tuberculosis* during THP-1 human macrophage infection. *Infect Immun* 2008;**76**(2):717-725.
- \*60. Tailleux L, Waddell SJ, Pelizzola M, et al. Probing host pathogen cross-talk by transcriptional profiling of both *Mycobacterium tuberculosis* and infected human dendritic cells and macrophages. *PLoS ONE*. 2008;**3**(1):e1403.  
This reference dissects the interaction of *Mycobacterium tuberculosis* with phagocytes through investigating the transcriptomes of both host cell and pathogen.
61. Storrie B, Desjardins M. The biogenesis of lysosomes: is it a kiss and run, continuous fusion and fission process? *Bioessays* 1996;**18**(11):895-903.
62. Duclos S, Corsini R, Desjardins M. Remodeling of endosomes during lysosome biogenesis involves 'kiss and run' fusion events regulated by rab5. *J Cell Sci* 2003;**116**(5):907-918.
63. Clemens DL. Characterization of the *Mycobacterium tuberculosis* phagosome. *Trends Microbiol* 1996;**4**(3):113-118.
64. Duclos S, Desjardins M. Subversion of a young phagosome: the survival strategies of intracellular pathogens. *Cell Microbiol* 2000;**2**(5):365-377.
65. Armstrong JA, Hart PD. Response of cultured macrophages to *Mycobacterium tuberculosis*, with observations on fusion of lysosomes with phagosomes. *J. Exp. Med.* 1971;**134**(3):713-740.
66. Russell DG, Dant J, Sturgillkoszycki S. *Mycobacterium avium*- and *Mycobacterium tuberculosis*-containing vacuoles are dynamic, fusion-competent vesicles that are accessible to glycosphingolipids from the host cell plasmalemma. *J Immunol* 1996;**156**(12):4764-4773.
67. Clemens DL, Horwitz MA. The *Mycobacterium tuberculosis* phagosome interacts with early endosomes and is accessible to exogenously administered transferrin. *J Exp Med* 1996;**184**(4):1349-1355.
68. Via LE, Deretic D, Ulmer RJ, Hibler NS, Huber LA, Deretic V. Arrest of mycobacterial phagosome maturation is caused by a block in vesicle fusion between stages controlled by rab5 and rab7. *J Biol Chem* 1997;**272**(20):13326-13331.
69. Xu SM, Cooper A, Sturgillkoszycki S, et al. Intracellular trafficking in *Mycobacterium tuberculosis* and *Mycobacterium avium*-infected macrophages. *J Immunol* 1994;**153**(6):2568-2578.
70. Clemens DL, Horwitz MA. Characterization of the *Mycobacterium tuberculosis* phagosome and evidence that phagosomal maturation is inhibited. *J Exp Med* 1995;**181**(1):257-270.
71. Crowle AJ, Dahl R, Ross E, May MH. Evidence that vesicles containing living, virulent *Mycobacterium tuberculosis* or *Mycobacterium avium* in cultured human macrophages are not acidic. *Infect Immun* 1991;**59**(5):1823-1831.
72. Oh YK, Straubinger RM. Intracellular fate of *Mycobacterium avium*: use of dual-label spectrofluorometry to investigate the influence of bacterial viability and opsonization on phagosomal pH and phagosome-lysosome interaction. *Infect Immun* 1996;**64**(1):319-325.
- \*73. Sturgillkoszycki S, Schlesinger PH, Chakraborty P, et al. Lack of acidification in *Mycobacterium* phagosomes produced by exclusion of the vesicular proton-ATPase. *Science* 1994;**263**(5147):678-681.  
This reference was the first to describe a mechanistic explanation for the lack of acidification of *Mycobacterium tuberculosis* phagosomes.

74. Beatty WL, Rhoades ER, Hsu DK, Liu FT, Russell DG. Association of a macrophage galactoside-binding protein with Mycobacterium-containing phagosomes. *Cell Microbiol* 2002;**4**(3):167-176.
  75. Ferrari G, Langen H, Naito M, Pieters J. A coat protein on phagosomes involved in the intracellular survival of mycobacteria. *Cell* 1999;**97**(4):435-447.
  76. Fratti RA, Chua J, Deretic V. Cellubrevin alterations and *Mycobacterium tuberculosis* phagosome maturation arrest. *J Biol Chem* 2002;**277**(19):17320-17326.
  77. Fratti RA, Chua J, Vergne I, Deretic V. *Mycobacterium tuberculosis* glycosylated phosphatidylinositol causes phagosome maturation arrest. *Proc Natl Acad Sci U S A* 2003;**100**(9):5437-5442.
  78. Welin A, Winberg ME, Abdalla H, et al. Incorporation of *Mycobacterium tuberculosis* lipoarabinomannan into macrophage membrane rafts is a prerequisite for the phagosomal maturation block. *Infect Immun*. 2008;**76**(7):2882-2887.
  79. Indrigo J, Hunter RLJ, Actor JK. Cord factor trehalose 6,6'-dimycolate (TDM) mediates trafficking events during mycobacterial infection of murine macrophages. *Microbiology* 2003;**149**(Pt 8):2049-2059.
  80. Malik ZA, Iyer SS, Kusner DJ. *Mycobacterium tuberculosis* phagosomes exhibit altered calmodulin-dependent signal transduction: Contribution to inhibition of phagosome-lysosome fusion and intracellular survival in human macrophages. *J Immunol* 2001;**166**(5):3392-3401.
  81. Malik ZA, Thompson CR, Hashimi S, Porter B, Iyer SS, Kusner DJ. *Mycobacterium tuberculosis* Blocks Ca<sup>2+</sup> Signaling and Phagosome Maturation in Human Macrophages Via Specific Inhibition of Sphingosine Kinase. *J Immunol* 2003;**170**(6):2811-2815.
  - \*82. Pethe K, Swenson DL, Alonso S, Anderson J, Wang C, Russell DG. Isolation of *Mycobacterium tuberculosis* mutants defective in the arrest of phagosome maturation. *Proc Natl Acad Sci U S A* 2004;**101**(37):13642-13647.
  - \*83. MacGurn JA, Cox JS. A genetic screen for *Mycobacterium tuberculosis* mutants defective for phagosome maturation arrest identifies components of the ESX-1 secretion system. *Infect Immun*. 2007;**75**(6):2668-2678.
- These two references identify mutants of *Mycobacterium tuberculosis* that are unable to arrest phagosome development.
84. Ganguly N, Siddiqui I, Sharma P. Role of *M. tuberculosis* RD-1 region encoded secretory proteins in protective response and virulence. *Tuberculosis* 2008;**88**(6):510-517.
  85. Greco E, De Spirito M, Papi M, Fossati M, Auricchio G, Fraziano M. CpG oligodeoxynucleotides induce Ca<sup>2+</sup>-dependent phospholipase D activity leading to phagolysosome maturation and intracellular mycobacterial growth inhibition in monocytes. *Biochem Biophys Res Commun*. 2006;**347**(4):963-969.
  86. Pieters J. *Mycobacterium tuberculosis* and the macrophage: maintaining a balance. *Cell Host Microb* 2008;**3**(6):399-407.
  87. Roberts EA, Deretic V. The *Mycobacterium tuberculosis* phagosome. *Methods Mol Biol*. 2008;**445**:439-449.
  88. Hampshire T, Soneji S, Bacon J, et al. Stationary phase gene expression of *Mycobacterium tuberculosis* following a progressive nutrient depletion: a model for persistent organisms? *Tuberculosis* 2004;**84**(3-4):228-238.

89. Betts JC, Lukey PT, Robb LC, McAdam RA, Duncan K. Evaluation of a nutrient starvation model of *Mycobacterium tuberculosis* persistence by gene and protein expression profiling. *Mol Microbiol* 2002;**43**(3):717-731.
90. Gold B, Rodriguez GM, Marras SA, Pentecost M, Smith I. The *Mycobacterium tuberculosis* IdeR is a dual functional regulator that controls transcription of genes involved in iron acquisition, iron storage and survival in macrophages. *Mol Microbiol.* 2001;**42**(3):851-865.
91. Rodriguez GM, Voskuil MI, Gold B, Schoolnik GK, Smith I. ideR, an Essential Gene in *Mycobacterium tuberculosis*: Role of IdeR in Iron-Dependent Gene Expression, Iron Metabolism, and Oxidative Stress Response. *Infect Immun* 2002;**70**(7):3371-3381.
92. Fontan PA, Aris V, Alvarez ME, et al. *Mycobacterium tuberculosis* sigma factor E regulon modulates the host inflammatory response. *J Infect Dis.* 2008;**198**(6):877-885.
93. Fisher MA, Plikaytis BB, Shinnick TM. Microarray analysis of the *Mycobacterium tuberculosis* transcriptional response to the acidic conditions found in phagosomes. *J Bacteriol* 2002;**184**(14):4025-4032.
94. Saviola B, Woolwine SC, Bishai WR. Isolation of acid-inducible genes of *Mycobacterium tuberculosis* with the use of recombinase-based *in vivo* expression technology. *Infect. Immun.* 2003;**71**(3):1379-1388.
95. Sherman DR, Voskuil M, Schnappinger D, Liao RL, Harrell MI, Schoolnik GK. Regulation of the *Mycobacterium tuberculosis* hypoxic response gene encoding alpha-crystallin. *Proc Natl Acad Sci U S A* 2001;**98**(13):7534-7539.
96. Park HD, Guinn KM, Harrell MI, et al. Rv3133c/dosR is a transcription factor that mediates the hypoxic response of *Mycobacterium tuberculosis*. *Mol Microbiol* 2003;**48**(3):833-843.
97. Bacon J, James BW, Wernisch L, et al. The influence of reduced oxygen availability on pathogenicity and gene expression in *Mycobacterium tuberculosis*. *Tuberculosis* 2004;**84**(3-4):205-217.
98. Muttucumaru DG, Roberts G, Hinds J, Stabler RA, Parish T. Gene expression profile of *Mycobacterium tuberculosis* in a non-replicating state. *Tuberculosis* 2004;**84**(3-4):239-246.
- \*\*99. Rustad TR, Harrell MI, Liao R, Sherman DR. The enduring hypoxic response of *Mycobacterium tuberculosis*. *PLoS ONE.* 2008;**3**(1):e1502.
- This reference demonstrates that the DevR(DosR) regulon is a transient response to hypoxia that is followed by an enduring hypoxic response regulon.
100. Wayne LG, Hayes LG. An in vitro model for sequential study of shutdown of *Mycobacterium tuberculosis* through two stages of nonreplicating persistence. *Infection and Immunity* 1996;**64**(6):2062-2069.
- \*101. Balazsi G, Heath AP, Shi L, Gennaro ML. The temporal response of the *Mycobacterium tuberculosis* gene regulatory network during growth arrest. *Mol Syst Biol.* 2008;**4**:225.
- This reference characterizes the temporal response of the *Mycobacterium tuberculosis* transcriptional-regulatory network during the shift to dormancy.
102. Serbina NV, Jia T, Hohl TM, Pamer EG. Monocyte-mediated defense against microbial pathogens. *Annu Rev Immunol.* 2008;**26**:421-52.
103. Hemsworth GR, Kochan I. Secretion of antimycobacterial fatty acids by normal and activated macrophages. *Infect Immun* 1978;**19**(1):170-177.

104. Ogata K, Linzer BA, Zuberi RI, Ganz T, Lehrer RI, Catanzaro A. Activity of defensins from human neutrophilic granulocytes against *Mycobacterium avium-Mycobacterium intracellulare*. *Infect Immun* 1992;**60**(11):4720-4725.
  105. Fairbairn IP, Stober CB, Kumararatne DS, Lammas DA. ATP-mediated killing of intracellular mycobacteria by macrophages is a P2X(7)-dependent process inducing bacterial death by phagosome-lysosome fusion. *J Immunol* 2001;**167**(6):3300-3307.
  106. Gutierrez MG, Master SS, Singh SB, Taylor GA, Colombo MI, Deretic V. Autophagy is a defense mechanism inhibiting BCG and *Mycobacterium tuberculosis* survival in infected macrophages. *Cell* 2004;**119**(6):753-766.
  107. Biswas D, Qureshi OS, Lee WY, Croudace JE, Mura M, Lammas DA. ATP-induced autophagy is associated with rapid killing of intracellular mycobacteria within human monocytes/macrophages. *BMC Immunol.* 2008;**9**(1):e35.
  108. Venketaraman V, Dayaram YK, Talaue MT, Connell ND. Glutathione and nitrosoglutathione in macrophage defense against *Mycobacterium tuberculosis*. *Infect. Immun.* 2005;**73**(3):1886-1889.
  109. MacMicking JD, Taylor GA, McKinney JD. Immune control of tuberculosis by IFN- $\gamma$ -Inducible LRG-47. *Science* 2003;**302**(5645):654-659.
  110. Liu PT, Stenger S, Tang DH, Modlin RL. Cutting Edge: Vitamin D-mediated human antimicrobial activity against *Mycobacterium tuberculosis* is dependent on the induction of cathelicidin. *J Immunol* 2007;**179**(4):2060-2063.
  111. Koo IC, Ohol YM, Wu P, Morisaki JH, Cox JS, Brown EJ. Role for lysosomal enzyme beta-hexosaminidase in the control of mycobacteria infection. *Proc Natl Acad Sci U S A.* 2008;**105**(2):710-715.
  112. Armstrong JA, Hart PD. Phagolysosome-lysosome interactions in cultured macrophages infected with virulent tubercle bacilli: Reversal of the usual nonfusion pattern and observations on bacterial survival. *J Exp Med* 1975;**142**:1-16.
  113. Kusner DJ, Barton JA. ATP stimulates human macrophages to kill intracellular virulent *Mycobacterium tuberculosis* via calcium-dependent phagosome-lysosome fusion. *J Immunol* 2001;**167**(6):3308-3315.
  114. Anes E, Kuhnel MP, Bos E, Moniz-Pereira J, Habermann A, Griffiths G. Selected lipids activate phagosome actin assembly and maturation resulting in killing of pathogenic mycobacteria. *Nat Cell Biol* 2003;**5**(9):793-802.
  115. Deretic V, Singh S, Master S, et al. *Mycobacterium tuberculosis* inhibition of phagolysosome biogenesis and autophagy as a host defence mechanism. *Cell Microbiol* 2006;**8**(5):719-727.
  - \*116. Nathan C, Shiloh MU. Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. *Proc Natl Acad Sci U S A* 2000;**97**(16):8841-8848.
- This review summarizes the role of reactive oxygen and nitrogen intermediates in mammalian immunity.
117. Bogdan CT, Rollinghoff M, Diefenbach A. The role of nitric oxide in innate immunity. *Immunol Rev* 2000;**173**:17-26.
  118. Zahrt TC, Deretic V. Reactive nitrogen and oxygen intermediates and bacterial defenses: Unusual adaptations in *Mycobacterium tuberculosis*. *Antioxid Redox Signal* 2002;**4**(1):141-159.

- 1 119. Fang FC. Antimicrobial reactive oxygen and nitrogen species: concepts and controversies.  
2 *Nat Rev Microbiol.* 2004;**2**(10):820-832.
- 3 120. Balcewicz-Sablinska MK, Keane J, Kornfeld H, Remold HG. Pathogenic *Mycobacterium*  
4 *tuberculosis* evades apoptosis of host macrophages by release of TNF-R2, resulting in  
5 inactivation of TNF $\alpha$ ". *J Immunol* 1998;**161**(5):2636-2641.
- 6 121. Keane J, Shurtleff B, Kornfeld H. TNF-dependent BALB/c murine macrophage apoptosis  
7 following *Mycobacterium tuberculosis* infection inhibits bacillary growth in an IFN-  
8 gamma independent manner. *Tuberculosis* 2002;**82**(2-3):55-61.
- 9 122. Pan H, Yan BS, Rojas M, et al. Ipr1 gene mediates innate immunity to tuberculosis. *Nature.*  
10 2005;**434**(7034):767-772.
- 11 \*123. Randhawa AK, Ziltener HJ, Stokes RW. CD43 controls the intracellular growth of  
12 *Mycobacterium tuberculosis* through the induction of TNF-alpha-mediated apoptosis.  
13 *Cell Microbiol.* 2008;**10**(10):2105-2117.
- 14 This reference identifies a receptor-mediated mechanism for the induction of apoptosis of  
15 *Mycobacterium tuberculosis* infected macrophages.  
16
- 17 124. Miller RA, Britigan BE. The formation and biological significance of phagocyte derived  
18 oxidants. *J Invest Med* 1995;**43**(1):39-49.
- 19 125. O'Brien L, Roberts B, Andrew PW. *In vitro* interaction of *Mycobacterium tuberculosis* and  
20 macrophages: activation of anti-mycobacterial activity of macrophages and mechanisms  
21 of anti-mycobacterial activity. *Curr Top Microbiol Immunol.* 1996;**215**:97-130.
- 22 126. Manca C, Paul S, Barry CE, Freedman VH, Kaplan G. *Mycobacterium tuberculosis* catalase  
23 and peroxidase activities and resistance to oxidative killing in human monocytes in vitro.  
24 *Infect Immun* 1999;**67**(1):74-79.
- 25 127. Adams LB, Dinauer MC, Morgenstern DE, Krahenbuhl JL. Comparison of the roles of  
26 reactive oxygen and nitrogen intermediates in the host response to *Mycobacterium*  
27 *tuberculosis* using transgenic mice. *Tuber Lung Dis* 1997;**78**(5-6):237-246.
- 28 128. Cooper AM, Segal BH, Frank AA, Holland SM, Orme IM. Transient loss of resistance to  
29 pulmonary tuberculosis in p47(phox-/-) Mice. *Infect Immun* 2000;**68**(3):1231-1234.
- 30 129. Ng VH, Cox JS, Sousa AO, MacMicking JD, McKinney JD. Role of KatG catalase-  
31 peroxidase in mycobacterial pathogenesis: countering the phagocyte oxidative burst. *Mol*  
32 *Microbiol* 2004;**52**(5):1291-1302.
- 33 130. Rhoades ER, Orme IM. Susceptibility of a panel of virulent strains of *Mycobacterium*  
34 *tuberculosis* to reactive nitrogen intermediates. *Infect Immun* 1997;**65**(4):1189-1195.
- 35 131. Firmani MA, Riley LW. Reactive nitrogen intermediates have a bacteriostatic effect on  
36 *Mycobacterium tuberculosis* in vitro. *J. Clin. Microbiol.* 2002;**40**(9):3162-3166.
- 37 132. MacMicking JD, North RJ, LaCourse R, Mudgett JS, Shah SK, Nathan CF. Identification of  
38 nitric oxide synthase as a protective locus against tuberculosis. *Proc Natl Acad Sci U S A*  
39 1997;**94**(10):5243-5248.
- 40 133. Scanga CA, Mohan VP, Tanaka K, Alland D, Flynn JL, Chan J. The inducible nitric oxide  
41 synthase locus confers protection against aerogenic challenge of both clinical and  
42 laboratory strains of *Mycobacterium tuberculosis* in mice. *Infect Immun*  
43 2001;**69**(12):7711-7717.
- 44 134. Hisert KB, Kirksey MA, Gomez JE, et al. Identification of *Mycobacterium tuberculosis*  
45 counterimmune (cim) mutants in immunodeficient mice by differential screening. *Infect*  
46 *Immun* 2004;**72**(9):5315-5321.

- 1 135. Shiloh MU, MacMicking JD, Nicholson S, et al. Phenotype of mice and macrophages  
2 deficient in both phagocyte oxidase and inducible nitric oxide synthase. *Immunity*  
3 1999;**10**(1):29-38.
- 4 136. Miller BH, Fratti RA, Poschet JF, et al. Mycobacteria inhibit nitric oxide synthase  
5 recruitment to phagosomes during macrophage infection. *Infect. Immun.*  
6 2004;**72**(5):2872-2878.
- 7 137. Rooyackers AW, Stokes RW. Absence of complement receptor 3 results in reduced binding  
8 and ingestion of *Mycobacterium tuberculosis* but has no significant effect on the  
9 induction of reactive oxygen and nitrogen intermediates or on the survival of the bacteria  
10 in resident and interferon-gamma activated macrophages. *Microb Pathog* 2005;**39**(3):57-  
11 67.
- 12 138. Schneemann M, Schoedon G, Hofer S, Blau N, Guerrero L, Schaffner A. Nitric oxide  
13 synthase is not a constituent of the antimicrobial armature of human mononuclear  
14 phagocytes. *J Infect Dis* 1993;**167**:1358-1363.
- 15 139. Denis M. Human monocytes/macrophages: NO or no NO? *J Leukoc Biol* 1994;**55**:682-684.
- 16 140. Albina JE. On the expression of nitric oxide synthase by human macrophages. Why no NO?  
17 *J Leukoc Biol* 1995;**58**:643-649.
- 18 141. Condoneto A, Muscara MN, Grumach AS, Carneirosampaio MMS, Denucci G.  
19 Neutrophils and mononuclear cells from patients with Chronic Granulomatous Disease  
20 release Nitric Oxide. *Br J Clin Pharmacol* 1993;**35**(5):485-490.
- 21 142. Nicholson S, Bonecini-Almeida M-G, Lapa e Silva JR, et al. Inducible nitric oxide synthase  
22 in pulmonary alveolar macrophages from patients with tuberculosis. *J Exp Med*  
23 1996;**183**:2293-2302.
- 24 143. Nozaki Y, Hasegawa Y, Ichiyama S, Nakashima I, Shimokata K. Mechanism of nitric  
25 oxide-dependent killing of *Mycobacterium bovis* BCG in human alveolar macrophages.  
26 *Infect Immun* 1997;**65**(9):3644-3647.
- 27 144. Dlugovitzky D, Bay ML, Ratani L, et al. Influence of disease severity on nitrite and  
28 cytokine production by peripheral blood mononuclear cells (PBMC) from patients with  
29 pulmonary tuberculosis (TB). *Clin Exp Immunol.* 2000;**122**(3):343-349.
- 30 145. Thomassen MJ, Kavuru MS. Human alveolar macrophages and monocytes as a source and  
31 target for nitric oxide. *Int Immunopharmacol.* 2001;**1**(8):1479-1490.
- 32 146. Chang HY, Yang X. Proteases for cell suicide: functions and regulation of caspases.  
33 *Microbiol Mol Biol Rev* 2000;**64**(4):821-846.
- 34 147. Riendeau CJ, Kornfeld H. THP-1 cell apoptosis in response to Mycobacterial infection.  
35 *Infect. Immun.* 2003;**71**(1):254-259.
- 36 148. Lee J, Remold HG, Jeong MH, Kornfeld H. Macrophage apoptosis in response to high  
37 intracellular burden of *Mycobacterium tuberculosis* is mediated by a novel caspase-  
38 independent pathway. *J Immunol.* 2006;**176**(7):4267-474.
- 39 149. Keane J, Balcewicz-Sablinska MK, Remold HG, et al. Infection by *Mycobacterium*  
40 *tuberculosis* promotes human alveolar macrophage apoptosis. *Infect Immun*  
41 1997;**65**(1):298-304.
- 42 150. Fratazzi C, Arbeit RD, Carini C, et al. Macrophage apoptosis in mycobacterial infections. *J*  
43 *Leukoc Biol* 1999;**66**(5):763-764.
- 44 151. Oddo M, Renno T, Attinger A, Bakker T, Macdonald HR, Meylan PRA. Fas ligand-induced  
45 apoptosis of infected human macrophages reduces the viability of intracellular  
46 *Mycobacterium tuberculosis*. *J Immunol* 1998;**160**(11):5448-5454.



- 1 152. Duan L, Gan H, Golan DE, Remold HG. Critical role of mitochondrial damage in  
2 determining outcome of macrophage infection with *Mycobacterium tuberculosis*. *J*  
3 *Immunol* 2002;**169**(9):5181-5187.
- 4 \*153. Keane J, Remold HG, Kornfeld H. Virulent *Mycobacterium tuberculosis* strains evade  
5 apoptosis of infected alveolar macrophages. *J Immunol* 2000;**164**(4):2016-2020.  
6 This reference described for the first time how avirulent and virulent strains of *Mycobacterium*  
7 *tuberculosis* differed in their propensity to trigger apoptosis of infected macrophages.  
8
- 9 154. Gao LY, Abu Kwaik Y. Hijacking of apoptotic pathways by bacterial pathogens. *Microb*  
10 *Infect* 2000;**2**(14):1705-1719.
- 11 155. Sly LM, Hingley-Wilson SM, Reiner NE, McMaster WR. Survival of *Mycobacterium*  
12 *tuberculosis* in host macrophages involves resistance to apoptosis dependent upon  
13 induction of antiapoptotic Bcl-2 family member Mcl-1. *J Immunol* 2003;**170**(1):430-437.
- 14 \*156. Velmurugan K, Chen B, Miller JL, et al. *Mycobacterium tuberculosis nuoG* is a virulence  
15 gene that inhibits apoptosis of infected host cells. *PLoS Pathog* 2007;**3**(7):e110 epub.  
16 This reference identifies a gene that is required for *Mycobacterium tuberculosis* to inhibit  
17 macrophage apoptosis.  
18
- 19 157. Hinchey J, Lee S, Jeon BY, et al. Enhanced priming of adaptive immunity by a proapoptotic  
20 mutant of *Mycobacterium tuberculosis*. *J Clin Invest*. 2007;**117**(8):2279-2288.
- 21 158. Kendall SL, Movahedzadeh F, Rison SC, et al. The *Mycobacterium tuberculosis* dosRS two-  
22 component system is induced by multiple stresses. *Tuberculosis* 2004;**84**(3-4):247-255.
- 23 159. Hu Y, Kendall S, Stoker NG, Coates AR. The *Mycobacterium tuberculosis sigJ* gene  
24 controls sensitivity of the bacterium to hydrogen peroxide. *FEMS Microbiol Lett*  
25 2004;**237**(2):415-423.
- 26 160. Ohno H, Zhu G, Mohan VP, et al. The effects of reactive nitrogen intermediates on gene  
27 expression in *Mycobacterium tuberculosis*. *Cell Microbiol*, 2003: 637-648.
- 28 161. Voskuil MI, Schnappinger D, Visconti KC, et al. Inhibition of respiration by nitric oxide  
29 induces a *Mycobacterium tuberculosis* dormancy program. *J Exp Med* 2003;**198**(5):705-  
30 713.
- 31 162. Gill WP, Harik NS, Whiddon MR, Liao RP, Mittler JE, Sherman DR. A replication clock  
32 for *Mycobacterium tuberculosis*. *Nat Med*. 2009.
- 33 163. Turner J, D'Souza CD, Pearl JE, et al. CD8- and CD95/95L-dependent mechanisms of  
34 resistance in mice with chronic pulmonary tuberculosis. *Am J Resp Cell Mol Biol*  
35 2001;**24**(2):203-209.
- 36 164. Brookes RH, Pathan AA, McShane H, Hensmann M, Price DA, Hill AV. CD8+ T cell-  
37 mediated suppression of intracellular *Mycobacterium tuberculosis* growth in activated  
38 human macrophages. *Eur J Immunol*, 2003: 3293-3302.
- 39 165. D'Souza CD, Cooper AM, Frank AA, et al. A novel nonclassic beta 2-microglobulin-  
40 restricted mechanism influencing early lymphocyte accumulation and subsequent  
41 resistance to tuberculosis in the lung. *Am J Resp Cell Mol Biol* 2000;**23**(2):188-193.
- 42 166. Mogues T, Goodrich ME, Ryan L, LaCourse R, North RJ. The relative importance of T cell  
43 subsets in immunity and immunopathology of airborne *Mycobacterium tuberculosis*  
44 infection in mice. *J Exp Med* 2001;**193**(3):271-280.
- 45 167. Flynn JL, Chan J. Immunology of tuberculosis. *Annu Rev Immunol* 2001;**19**:93-129.
- 46 168. Orme I. Adaptive immunity to mycobacteria. *Curr Opin Microbiol* 2004;**7**(1):58-61.

169. Cooper AM. Cell-mediated immune responses in tuberculosis. *Annu Rev Immunol.* 2009 **27**:393-422.
170. Schaible UE, Sturgillkoszycki S, Schlesinger PH, Russell DG. Cytokine activation leads to acidification and increases maturation of *Mycobacterium avium*-containing phagosomes in murine macrophages. *J Immunol* 1998;**160**(3):1290-1296.
171. Laochumroonvorapong P, Paul S, Elkon KB, Kaplan G. H<sub>2</sub>O<sub>2</sub> induces monocyte apoptosis and reduces viability of *Mycobacterium avium*-*M. intracellulare* within cultured human monocytes. *Infect Immun* 1996;**64**(2):452-459.
172. Reed MB, Domenech P, Manca C, et al. A glycolipid of hypervirulent tuberculosis strains that inhibits the innate immune response. *Nature* 2004 **431**(7004):84-87.
173. Pathak SK, Basu S, Bhattacharyya A, Pathak S, Kundu M, Basu J. *Mycobacterium tuberculosis* lipoarabinomannan-mediated IRAK-M induction negatively regulates Toll-like receptor-dependent interleukin-12 p40 production in macrophages. *J Biol Chem.* 2005;**280**(52):42794-42800.
174. Dao DN, Sweeney K, Hsu T, et al. Mycolic acid modification by the mmaA4 gene of *M. tuberculosis* modulates IL-12 production. *PLoS Pathog.* 2008;**4**(6):e1000081-e1000081.
175. Fortune SM, Solache A, Jaeger A, et al. *Mycobacterium tuberculosis* Inhibits Macrophage Responses to IFN- $\gamma$  through Myeloid Differentiation Factor 88-Dependent and -Independent Mechanisms. *J Immunol* 2004;**172**(10):6272-6280.
176. Chan J, Xing Y, Magliozzo RS, Bloom BR. Killing of virulent *Mycobacterium tuberculosis* by reactive nitrogen intermediates produced by activated murine macrophages. *J Exp Med* 1992;**175**(4):1111-1122.
177. McKinney JD, zu Bentrup KH, Munoz-Elias EJ, et al. Persistence of *Mycobacterium tuberculosis* in macrophages and mice requires the glyoxylate shunt enzyme isocitrate lyase. *Nature.* 2000;**406**(6797):735-738.
178. Shi L, Sohaskey CD, Kana BD, et al. Changes in energy metabolism of *Mycobacterium tuberculosis* in mouse lung and under *in vitro* conditions affecting aerobic respiration. *Proc Natl Acad Sci U S A.* 2005;**102**(43):15629-15634.
179. Murphy DJ, Brown JR. Identification of gene targets against dormant phase *Mycobacterium tuberculosis* infections. *BMC Infect Dis.* 2007; **7**:84.
180. Voskuil MI, Visconti KC, Schoolnik GK. *Mycobacterium tuberculosis* gene expression during adaptation to stationary phase and low-oxygen dormancy. *Tuberculosis* 2004;**84**(3-4):218-227.
181. Lewis KN, Liao R, Guinn KM, et al. Deletion of RD1 from *Mycobacterium tuberculosis* mimics bacille Calmette-Guerin attenuation. *J Infect Dis* 2003;**187**(1):117-123.
- \*182. Golby P, Hatch KA, Bacon J, et al. Comparative transcriptomics reveals key gene expression differences between the human and bovine pathogens of the *Mycobacterium tuberculosis* complex. *Microbiology.* 2007;**153**(Pt 10):3323-3336.
- This reference identifies significant differences between *Mycobacterium tuberculosis* and *M. bovis* gene expression, illustrating the problem with using *M. bovis* BCG as a model for *M. tuberculosis*.
183. Flesch IEA, Kaufmann SHE. Mechanisms involved in mycobacterial growth inhibition by gamma interferon-activated bone marrow macrophages: role of reactive nitrogen intermediates. *Infect Immun* 1991;**59**(1):3213-3218.

184. Shi L, Jung YJ, Tyagi S, Gennaro ML, North RJ. Expression of Th1-mediated immunity in mouse lungs induces a *Mycobacterium tuberculosis* transcription pattern characteristic of nonreplicating persistence. *Proc Natl Acad Sci U S A* 2003;**100**(1):241-246.
  185. Dubnau E, Chan J, Mohan VP, Smith I. Responses of *Mycobacterium tuberculosis* to growth in the mouse lung. *Infect. Immun.* 2005;**73**(6):3754-3757.
  186. Talaat AM, Ward SK, Wu CW, et al. Mycobacterial bacilli are metabolically active during chronic Tuberculosis in murine lungs: Insights from genome-wide transcriptional profiling. *J Bacteriol.* 2007;**189**(11):4265-4274.
  187. Shi L, Sohaskey CD, North RJ, Gennaro ML. Transcriptional characterization of the antioxidant response of *Mycobacterium tuberculosis* *in vivo* and during adaptation to hypoxia *in vitro*. *Tuberculosis* 2008;**88**(1):1-6.
  188. Fenhalls G, Stevens L, Moses L, et al. *In situ* detection of *Mycobacterium tuberculosis* transcripts in human lung granulomas reveals differential gene expression in necrotic lesions. *Infect Immun* 2002;**70**(11):6330-6338.
  189. Garton NJ, Waddell SJ, Sherratt AL, et al. Cytological and transcript analyses reveal fat and lazy persister-like bacilli in tuberculous sputum. *PLoS Med.* 2008 **5**(4):e75.
  190. Capuano SV, III, Croix DA, Pawar S, et al. Experimental *Mycobacterium tuberculosis* Infection of Cynomolgus Macaques closely resembles the various manifestations of human *M. tuberculosis* Infection. *Infect Immun* 2003;**71**(10):5831-5844.
  191. Lin PL, Pawar S, Myers A, et al. Early events in *Mycobacterium tuberculosis* infection in Cynomolgus Macaques. *Infect Immun* 2006;**74**(7):3790-3803.
  192. Lee J, Hartman M, Kornfeld H. Macrophage apoptosis in tuberculosis. *Yonsei Med J.* 2009 **50**(1):1-11.
  193. Gan H, Lee J, Ren F, Chen M, Kornfeld H, Remold HG. *Mycobacterium tuberculosis* blocks crosslinking of annexin-1 and apoptotic envelope formation on infected macrophages to maintain virulence. *Nat Immunol.* 2008;**9**(10):1189- 1197.
  194. Spira A, Carroll JD, Liu G, et al. Apoptosis genes in human alveolar macrophages infected with virulent or attenuated *Mycobacterium tuberculosis*: a pivotal role for tumor necrosis factor. *Am J Respir Cell Mol Biol* 2003;**29**(5):545-551.
  195. Gao Q, Kripke K, Arinc Z, Voskuil M, Small P. Comparative expression studies of a complex phenotype: cord formation in *Mycobacterium tuberculosis*. *Tuberculosis* 2004;**84**(3-4):188-196.
  196. Walters SB, Dubnau E, Kolesnikova I, Laval F, Daffe M, Smith I. The *Mycobacterium tuberculosis* PhoPR two-component system regulates genes essential for virulence and complex lipid biosynthesis. *Mol Microbiol* 2006;**60**(2):312-330.
  - \*197. Lee JS, Krause R, Schreiber J, et al. Mutation in the transcriptional regulator PhoP contributes to avirulence of *Mycobacterium tuberculosis* H37Ra strain. *Cell Host Microb* 2008;**3**(2):97-103.
- This reference describes a major contributory factor in the avirulence of *Mycobacterium tuberculosis*, strain H37Ra when compared to the virulent strain H37Rv.
198. Hingley-Wilson SM, Sambandamurthy VK, Jacobs WRJ. Survival perspectives from the world's most successful pathogen, *Mycobacterium tuberculosis*. *Nat Immunol*, 2003: 949-955.
  199. Ward SK, Hoyer EA, Talaat AM. The global responses of *Mycobacterium tuberculosis* to physiological levels of copper. *J Bacteriol.* 2008;**190**(8):2939-2946.

- 1 200. Manganelli R, Voskuil MI, Schoolnik GK, Dubnau E, Gomez M, Smith I. Role of the  
2 extracytoplasmic-function sigma factor sigma(H) in *Mycobacterium tuberculosis* global  
3 gene expression. *Mol Microbiol.* 2002;**45**(2):365-374.
- 4 201. Manganelli R, Voskuil MI, Schoolnik GK, Smith I. The *Mycobacterium tuberculosis* ECF  
5 sigma factor sigmaE: role in global gene expression and survival in macrophages. *Mol*  
6 *Microbiol.* 2001;**41**(2):423-437.
- 7
- 8 301. WHO. Global tuberculosis control - epidemiology, strategy, financing. *Global Report*  
9 2009;**2009.411**:[http://www.who.int/tb/publications/global\\_report/2009/en/index.html](http://www.who.int/tb/publications/global_report/2009/en/index.html).

Figure 1




Figure 2




Figure 3

